



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/82, 9/88, 9/10, 9/00, 15/11, A01H 5/00	A2	(11) International Publication Number: WO 00/22150 (43) International Publication Date: 20 April 2000 (20.04.00)
--	----	--

(21) International Application Number: PCT/US99/23180

(22) International Filing Date: 5 October 1999 (05.10.99)

(30) Priority Data:
09/172,339 14 October 1998 (14.10.98) US(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application
US 09/172,339 (CON)
Filed on 14 October 1998 (14.10.98)

(71) Applicant (for all designated States except US): PIONEER HI-BRED INTERNATIONAL, INC. [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50319 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MEYER, Terry, EuClaire [US/US]; 4338 101st Street, Urbandale, IA 50322 (US). YALPANI, Nasser [CA/US]; 6041 North Winwood Drive, Johnston, IA 50131 (US).



(74) Agents: SPRUILL, W., Murray et al.; Alston & Bird LLP, P.O. Drawer 34009, Charlotte, NC 28234-4009 (US).

(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

Without international search report and to be republished upon receipt of that report.

(54) Title: LIMONENE AND OTHER DOWNSTREAM METABOLITES OF GERANYL PYROPHOSPHATE FOR INSECT CONTROL IN PLANTS



(57) Abstract

Methods for manipulating metabolic pathway in plants, particularly those pathways involved in the biosynthesis of monoterpenes are provided. Methods are directed at transforming plants with one or more nucleotide sequences encoding the enzyme GPP synthase, and the monoterpene synthases limonene-, carveol and S-linalool synthase. Methods for creating or enhancing resistance to insects in plants by transforming plants with GPP- and/or monoterpene synthases, to generate plants producing monoterpenes in amount effective for resistance to insects are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

LIMONENE AND OTHER DOWNSTREAM METABOLITES OF GERANYL PYROPHOSPHATE FOR INSECT CONTROL IN PLANTS

FIELD OF THE INVENTION

This invention relates to methods for genetic manipulation of metabolic pathways in plants, particularly to transforming plants with genes involved in monoterpene biosynthesis and resistance to insects.

5

BACKGROUND OF THE INVENTION

Numerous insects are serious pests of common agricultural crops. One method of controlling insects has been to apply insecticidal organic, semiorganic or organometallic chemicals to crops. This method has numerous, art-recognized environmental and public health problems. A more recent method of control of insect pests has been the use of biological control organisms which are typically natural predators of the troublesome insects. These include other insects such as trachonid wasps, fungi such as *Beauveria Bassiana*, and bacteria such as *Bacillus thuringiensis* cv., commonly referred to as "Bt". However, it is difficult to apply biological control organisms to large areas, and even more difficult to cause those living organisms to remain in the treated area for an extended period. Still more recently, techniques in recombinant DNA have provided the opportunity to insert into plant cells cloned genes which express insecticidal toxins derived from biological control organisms such as Bt. This technology has given rise to concerns about eventual insect resistance to well-known, naturally occurring insect toxins, particularly in the face of heavy selection pressure, which may occur in some areas. Thus, a continuing need exists to identify naturally occurring insecticidal toxins which can be formed by plant cells directly by expression of structural genes not normally present in the plant.

25 Southern Corn Rootworm (*Diabrotica undecimpunctata howardi* Barber) is a particularly difficult pest to control or eradicate. It attacks the plant below the soil line, where insecticides are difficult or impossible to apply effectively. In addition, it is resistant to a number of otherwise effective chemical and biological control agents, including Bt toxins and some lectins.

The monoterpene, limonene, 1-methyl-4 (1-methylethenyl) cyclohexene, p-mentha-1,8-diene (Entry No. 5371, Merck Index 11th Ed.), occurs naturally in various ethereal oils, particularly oils of lemon, orange, caraway, dill and bergamot. It is a valuable industrial chemical. Some limonene is prepared by
5 extraction from plants of the mint family, a large quantity is obtained from citrus oils, which are typically 80-90% limonene, and some is obtained from pine oil. It is also synthesized chemically and finds use as a solvent and cleaning agent (in the manufacture of synthetic pine oil), as an expectorant, as a wetting and dispersing agent, as a monomer in the manufacture of various polymeric resins, as a flavorant
10 and a precursor in the synthesis of the flavorant carvone, and as a polymerization inhibitor in storage of the tetrafluoroethylene monomer used in the manufacture of polytetrafluoroethylene (PTFE).

Geranyl diphosphate (GPP) synthase catalyzes the first committed step of monoterpene biosynthesis by the condensation of dimethylallyl diphosphate
15 (DMAPP) and isopentenyl diphosphate (IPP) to form GPP, the immediate acyclic precursor of monoterpenes. GPP is converted to (-)-4S-limonene by the catalytic action of (-)-4S-limonene synthase (cyclase). (-)-4S-limonene is converted to (-)-trans-carveol by the action of (-)-trans-carveol synthase, also referred to as limonene-6-hydroxylase. (-)-trans-carveol is converted to carvone by the action of
20 (-)-trans-carveol dehydrogenase, also referred to as carvone synthase. GPP is also converted to the monoterpene S-linalool by the action of S-linalool synthase. Thus, GPP is the precursor of (-)-4S-limonene and its downstream metabolites (-)-trans-carveol and carvone; as well as the precursor of S-linalool. See Figure 1; Wise *et al.* (1997) In "*Comprehensive Natural Products Chemistry: Isoprenoids*,
25 *Vol. 2*" (Cane, D.E., ed.), Elsevier Science, Oxford (in press); Gershenzon *et al.* (1989) *Plant Physiol.* 89:1351-1357; Pichersky *et al.* (1994) *Plant Physiol.* 106:1533-1540. Unlike the mechanistically-related prenyltransferases farnesyl diphosphate (FPP) synthase and geranylgeranyl diphosphate (GGPP) synthase, which produce GPP as intermediates and which are nearly ubiquitous (Ogura *et al.*
30 (1997) In "*Dynamic Aspects of Natural Products Chemistry*" (Ogura, K. and Sankawa, U., eds.), Kodansha/Harwood Academic Publishers, Tokyo, pp. 1-23), GPP synthase is largely restricted to plant species that produce abundant quantities of monoterpenes.

Because both farnesyl diphosphate synthase and geranylgeranyl diphosphate synthase produce only negligible levels of GPP as a free intermediate on route to FPP and GGPP (Ogura *et al.* (1997) In "*Dynamic Aspects of Natural Products Chemistry*" (Ogura, K. and Sankawa, U., eds.), Kodansha/Harwood Academic Publishers, Tokyo, pp. 1-23), it is geranyl diphosphate synthase that provides the crucial link between primary metabolism and monoterpene biosynthesis and that serves as the essential driver of monoterpene biosynthesis (Wise *et al.* (1997) In "*Comprehensive Natural Products Chemistry: Isoprenoids, Vol. 2*" (Cane, D.E., ed.), Elsevier Science, Oxford (in press)).

GPP synthase has been isolated from several plant sources, including grape, geranium, sage (Croteau *et al.* (1989) *Arch. Biochem. Biophys.* 271:524-535; Heide *et al.* (1989) *Arch. Biochem. Biophys.* 273:331-338; Suga *et al.* (1991) *Phytochemistry* 30:1757-1761; Clastre *et al.* (1993) *Plant Physiol.* 102:205-211); however, the enzyme has not been characterized in any detail, and only the enzyme from grape has been purified to homogeneity.

A cDNA encoding 4S-limonene synthase from oil glands of spearmint has been described in Colby *et al.* (1993) *J. Biol. Chem.* 268(31): 23016-23024 and is available in the GenbankTM/EMBL database and identified by the accession number L13459.

cDNAs encoding S-linalool synthase from *Clarkia Breweri* have been described in WO 97/15584, along with methods directed at using the cDNA for enhancing the flavor and smell of plants; and in Dudareva *et al.* (1996) *Plant Cell* 8 (7):1137-1148, also available in the GenbankTM/EMBL database and identified by the accession number 1491939.

To exploit recombinant methods to increase monoterpene yield in monoterpene-producing species, or to genetically engineer the monoterpene biosynthetic pathway into non-producing species, it would be highly beneficial to manipulate a GPP synthase gene. Accordingly, the invention relates to expressing GPP synthase in combination with selected monoterpene synthases such as (-)-limonene synthase, S-linalool synthase, and subsequent and related pathway enzymes for production of the corresponding monoterpene product(s).

SUMMARY OF THE INVENTION

The invention provides methods for manipulating metabolic pathways in plants, particularly those pathways that are involved in the biosynthesis of
5 monoterpenes. Methods are directed at transforming plants, plant tissues and cells with one or more nucleotide sequences encoding the enzyme GPP synthase, and the monoterpene synthases limonene-, carveol and S-linalool synthase.

Methods are also provided for creating or enhancing resistance to insects in plants by transforming plants with GPP- and/or monoterpene synthases, to generate
10 plants producing monoterpenes in amounts effective for resistance to insects.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the biosynthetic pathways involved in production of the monoterpenes limonene, carveol, and S-linalool.
15

Figure 2 schematically illustrates the plasmid construct comprising the ubiquitin promoter and a monoterpene synthesis pathway enzyme.

Figure 3 schematically illustrates the plasmid construct PHP14099 for
20 expression of GPP- and S-linalool synthases, and production of S-linalool in plants.

Figure 4 schematically illustrates the plasmid construct PHP14100 for
25 expression of GPP-, limonene- and carveol synthases; and production of limonene and carveol.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides methods for manipulating metabolic pathways in plant cells; particularly those pathways related to synthesis of
30 monoterpenes including but not limited to the monoterpenes (-)-4S-limonene (herein limonene), (-)-trans-carveol (herein carveol) and S-linalool in plants, plant cells and specific plant tissues. These monoterpenes have insecticidal and/or repellent activity against insect pests.

Accordingly, the invention provides methods useful for increasing monoterpene yield in monoterpene-producing species such as mint, and for producing monoterpenes in species which typically do not produce monoterpenes such as maize. The invention also provides methods for creating or enhancing
5 resistance to insects in plants by transforming plants with nucleotide sequences encoding monoterpene synthesis pathway enzymes and generating transformed plants which produce effective amounts of desired monoterpenes in the plant. In this aspect, by "effective amount" is intended that amount of a monoterpene, alone or in combination with other agents, that can effect a reduction, amelioration,
10 prevention, or elimination of a plant-insect interaction.

Thus, the methods of the invention are directed at transforming plant cells with at least one nucleotide sequence encoding a monoterpene synthesis pathway enzyme selected from GPP synthase, limonene synthase, carveol synthase, S-linalool synthase, or combinations thereof.

15 By "monoterpene synthase" is intended an enzyme which catalyzes a reaction having at least one monoterpene as the product of the reaction. More specifically, monoterpene synthases utilized in the methods of the invention include, but are not limited to limonene synthase, carveol synthase (limonene 6-hydroxylase) and S-linalool synthase.

20 By "transgenic plant" is meant any plant or plant cell that has become transformed by the introduction, stable and heritable incorporation, into the subject plant or plant cell, of either native DNA that is under the control of a promoter other than the promoter that typically drives expression of that DNA in a wild-type plant, and that has been introduced back into its host plant, or foreign DNA, i.e.
25 DNA encoding for a protein not normally found within that plant species.

"Plantlet" refers to a plant sufficiently developed to have a shoot and a root that is asexually reproduced by cell culture.

"Explant" refers to a section or piece of tissue from any part of a plant for culturing.

30 The term "callus" and its plural "calli", refer to an unorganized group of cells formed in response to cutting, severing, or other injury inflicted on plant tissue. Excised pieces of plant tissue and isolated cells can be induced to form callus under the appropriate culture conditions. Callus can be maintained in

culture for a considerable time by transferring or subculturing parts of the callus to fresh medium at regular intervals. The transfer of callus to liquid medium leads to dispersion of the tissue and the formation of a plant cell suspension culture. Callus can be induced to undergo organized development to form shoots and roots.

5 "Embryoid" refers to a structure similar in appearance to a plant zygotic embryo.

By the term "taxon" herein is meant a unit of botanical classification of genus or lower. It thus includes genus, species, cultivar, variety, variant, and other minor taxonomic groups that lack a consistent nomenclature.

10 "Somatic hybrid" and "somatic hybridization" refers generally to stable combination of cellular material, be it protoplast/protoplast or protoplast/cytoplasm combinations, and includes cybrids and cybridization.

A "replicon" is any genetic element (e.g., plasmid, chromosome, virus) that functions as a unit of DNA replication *in vivo*; i.e., capable of replication under its
15 own control.

As used herein, the term "nucleotide sequence" means a DNA or RNA sequence, and can include a cDNA, or genomic DNA, or synthetic DNA sequence, a structural gene or a fragment thereof, or an mRNA sequence, that encodes an active or functional polypeptide.

20 A "vector" is a replicon, such as a plasmid, phage, or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

A DNA "coding sequence" is a DNA sequence which is transcribed and translated into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a
25 start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but is not limited to, procaryotic sequences, cDNA from eucaryotic mRNA, genomic DNA sequences from eucaryotic (e.g., mammalian) DNA, and even synthetic DNA sequences. A
30 polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

A "promoter sequence" or a "promoter" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream

(3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bound at its 3' terminus by the translation start codon (ATG) of a coding sequence and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription. Within the
5 promoter sequence will be found a transcription initiation site, as well as protein binding domains responsible for the binding of RNA polymerase. Eucaryotic promoters will often, but not always, contain "TATA" boxes and "CAT" boxes.

DNA "control sequences" refers collectively to promoter sequences, ribosome binding sites, polyadenylation signals, transcription termination
10 sequences, upstream regulatory domains, enhancers, and the like, which collectively provide for the transcription and translation of a coding sequence in a host cell.

A coding sequence is "operably linked to" or "under the control of" control sequences in a cell when RNA polymerase will bind the promoter sequence and
15 transcribe the mRNA, which is then translated into the polypeptide encoded by the coding sequence.

A "host cell" is a cell which has been transformed, or is capable of undergoing transformation, by an exogenous DNA sequence.

A cell has been "transformed" by endogenous or exogenous DNA when
20 such DNA has been introduced inside the cell membrane. The DNA may or may not be integrated into (covalently linked to) chromosomal DNA making up the genome of the transformed cell. In procaryotes, for example, the DNA may be maintained on an episomal element, such as a plasmid. With respect to eucaryotic cells, a stably transformed cell is one in which the DNA has become integrated into
25 the chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eucaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the DNA.

A "clone" is a population of cells derived from a single cell or common
30 ancestor by mitosis. A "cell line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations.

A "heterologous" region of a DNA construct is an identifiable segment of DNA within or attached to another DNA molecule that is not found in association

with the other molecule in nature. Thus, when the heterologous region encodes a bacterial gene, the gene will usually be flanked by DNA that does not flank the bacterial gene in the genome of the source bacterium. Another example of a heterologous coding sequence is a construct where the coding sequence itself is not found in nature (e.g., synthetic sequences having codons different from the native gene). "Heterologous" DNA also refers to DNA not found within the host cell in nature. Allelic variation or naturally occurring mutational events do not give rise to a heterologous region of DNA, as these terms are used herein. "Native", "autologous" or "endogenous" DNA, as used herein, refer to DNA that is typically present in the host in nature.

The term "polypeptide" as used herein is used in its broadest sense, i.e., any polymer of amino acids (dipeptide or greater) linked through peptide bonds. Thus, the term "polypeptide" includes proteins, oligopeptides, protein fragments, analogues, muteins, fusion proteins and the like. The term also encompasses amino acid polymers as described above that include additional non-amino acid moieties. Thus, the term "polypeptide" includes glycoproteins, lipoproteins, phosphoproteins, metalloproteins, nucleoproteins, as well as other conjugated proteins. The term "polypeptide" contemplates polypeptides as defined above that are recombinantly produced, isolated from an appropriate source, or synthesized.

By the term "vector" herein is meant a DNA sequence which is able to replicate and express a foreign gene in a host cell. Typically, the vector has one or more endonuclease recognition sites which may be cut in a predictable fashion by use of the appropriate enzyme. Such vectors are preferably constructed to include additional structural gene sequences imparting antibiotic or herbicide resistance, which then serve as selectable markers to identify and separate transformed cells. Preferred selection agents include kanamycin, chlorosulfuron, phosphinothricin, hygromycin and methotrexate, and preferred markers are genes conferring resistance to these compounds. A cell in which the foreign genetic material in a vector is functionally expressed has been "transformed" by the vector and is referred to as a "transformant." A particularly preferred vector is a plasmid, by which is meant a circular double-stranded DNA molecule that is not a part of the chromosomes of the cell.

In carrying out this invention, it will be appreciated that numerous plant expression cassettes and vectors are well known in the art. By the term "expression cassette" is meant a complete set of control sequences including initiation, promoter and termination sequences which function in a plant cell when they flank a structural gene in the proper reading frame. Expression cassettes frequently and preferably contain an assortment of restriction sites suitable for cleavage and insertion of any desired structural gene. It is important that the cloned gene have a start codon in the correct reading frame for the structural sequence. In addition, the plant expression cassette preferably includes a strong promoter sequence at one end to cause the gene to be transcribed at a high frequency, and a poly-A recognition sequence at the other end for proper processing and transport of the messenger RNA. An example of such a preferred (empty) expression cassette into which the DNA sequence of the present invention can be inserted is the pPHI414 plasmid developed by Beach *et al.* of Pioneer Hi-Bred International, Inc., Johnston, IA. Highly preferred plant expression cassettes are designed to include one or more selectable marker genes, such as kanamycin resistance or herbicide tolerance genes.

Methods of the invention include those for manipulating a metabolic pathway in a plant cell by transforming with nucleotide sequences for native limonene-, GPP-, carveol- and S-linalool synthase genes, by transforming with nucleotide sequences encoding amino acid sequences for the respective proteins encoded thereby, as well as fragments and variants thereof. Such native sequences are set forth in SEQ ID NOs: 1-8. SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5 and SEQ ID NO:7 set forth the nucleotide sequences for limonene-, GPP-, carveol-, and S-linalool synthase respectively; the corresponding amino acid sequences set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, and SEQ ID NO:8 respectively. The methods encompass using the sequences or corresponding antisense sequences in modulating the expression of enzymes involved in monoterpene synthesis, as well as production of monoterpenes in a plant or plant cell. That is, the coding sequences are used to increase the expression of an enzyme while antisense sequences are used to decrease expression. In this aspect, blocking the expression of an enzyme within a pathway by antisense sequences can

be used to accumulate the substrates of that enzyme or to drive the pathway to another end product.

It is recognized that the methods of the invention could be used to manipulate metabolic pathways involving reactions downstream of those catalyzed by GPP- and a monoterpene synthase such as limonene-, carveol- and/or S-linalool synthase as described herein; by transforming plants with nucleotide sequences encoding GPP synthase and at least one monoterpene synthase, or antisense sequences thereof. In this aspect, the methods of the invention encompass manipulating the pathway involving production of the compounds including but not limited to (-)-carvone, (-)-trans-isopiperitol and (-)-trans-isopiperitinone.

Antisense RNA or DNA can be utilized for the accumulation of a particular monoterpene. Alternatively, homologous plant sequences or partial plant sequences can be used. For example, an antisense carveol synthase sequence can be used to cause the accumulation of limonene. In this manner, the metabolic pathway of interest can be manipulated for the high production of any particular monoterpene of interest in the pathway.

Likewise, the pathway can be manipulated to decrease levels of a particular compound by transformation of antisense sequences which prevent the conversion of the precursor compound into the particular compound being regulated. For example, conversion of GPP to S-linalool can be blocked by antisense sequences to S-linalool synthase.

In the same manner, to increase the biosynthesis of a particular desired monoterpene, antisense constructs can be used to block the conversion of a common substrate to one monoterpene, thereby shunting the common substrate to the pathway for the desired monoterpene, while additionally blocking downstream conversion of the desired monoterpene to a further downstream metabolite. For example, an antisense sequence to S-linalool synthase can be used to shunt the common substrate GPP to the pathway for limonene biosynthesis, while additionally antisense sequence to carveol synthase can be used to block conversion of limonene to carveol.

Any means for producing a plant comprising GPP- and at least one monoterpene synthase coding sequence are encompassed by the methods of the present invention. For example, the second (or additional) gene of interest can be

used to transform a plant at the same time as the GPP synthase gene (cotransformation); the second gene can be introduced into a plant that has already been transformed with the GPP synthase gene; GPP synthase can be transformed into a plant has already been transformed with the second gene; or alternatively, transformed plants, one expressing the GPP synthase and one expressing the second gene, can be crossed to bring the genes together in the same plant. Subsequent crosses or transformations can bring additional sequences together in the plant.

The use of fragments and variants of the nucleotide and amino acid sequences are encompassed within the scope of the invention. By "fragment" is intended a portion of the nucleotide sequence or a portion of the amino acid sequence and hence protein encoded thereby. Fragments of a nucleotide sequence may encode protein fragments that retain the biological activity of the native protein. Alternatively, fragments of a nucleotide sequence that are useful as hybridization probes generally do not encode fragment proteins retaining biological activity. Thus, fragments of a nucleotide sequence may range from at least about 15 nucleotides, about 50 nucleotides, about 100 nucleotides, and up to the entire nucleotide sequence encoding the monoterpene synthase proteins utilized in the methods of the invention. Fragments of the invention include antisense sequences used to decrease expression of the monoterpene synthase genes utilized in the methods of the invention. Such antisense fragments may vary in length ranging from at least about 15 nucleotides, about 50 nucleotides, about 100 nucleotides, up to and including the entire coding sequence.

By "variants" is intended substantially similar sequences. For nucleotide sequences, conservative variants include those sequences that, because of the degeneracy of the genetic code, encode the amino acid sequence of the GPP- and carveol synthase genes. Generally, nucleotide sequence variants of the invention will have at least 70%, generally, 80%, preferably up to 90-95% sequence identity to the native nucleotide sequence.

By "variant" protein is intended a protein derived from the native protein by deletion (so-called truncation) or addition of one or more amino acids to the N-terminal and/or C-terminal end of the native protein; deletion or addition of one or more amino acids at one or more sites in the native protein; or substitution of one

or more amino acids at one or more sites in the native protein. Such variants may result from, for example, genetic polymorphism or from human manipulation. Methods for such manipulations are generally known in the art.

For example, amino acid sequence variants of the polypeptide can be
5 prepared by mutations in the DNA sequence encoding the native protein of interest. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Walker and Gaastra, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York); Kunkel (1985) *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel *et al.* (1987) *Methods Enzymol.*
10 154:367-382; Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor, New York); U.S. Patent No. 4,873,192; and the references cited therein; herein incorporated by reference. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of Dayhoff *et al.* (1978) *Atlas of Protein Sequence and*
15 *Structure* (Natl. Biomed. Res. Found., Washington, D.C.), herein incorporated by reference. Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be preferred.

In constructing variants of the limonene-, GPP-, carveol and S-linalool synthase proteins of interest, modifications to the nucleotide sequences encoding
20 the variants will be made such that variants continue to possess the desired activity. Obviously, any mutations made in the DNA encoding the variant protein must not place the sequence out of reading frame and preferably will not create complementary regions that could produce excessive secondary mRNA structure. See EP Patent Application Publication No. 75,444.

25 Thus, nucleotide sequences utilized in the methods of the invention and the proteins encoded thereby include the native forms as well as variants thereof. The variant proteins will be substantially homologous and functionally equivalent to the native proteins. A variant of a native protein is "substantially homologous" to the native protein when at least about 80-85%, more preferably at least about 90%,
30 and most preferably at least about 95% of its amino acid sequence is identical to the amino acid sequence of the native protein. DNA sequences that are substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that particular system.

Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Maniatis *et al.* (1982) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York; Brown, T.A. *Gene Cloning: An Introduction* (2nd Ed.) Chapman & Hall, London (1990).

5 Alternatively, sequence identity can be determined utilizing a number of available computer programs including but not limited to CLUSTAL in the PC/Gene program (available from Intelligenetics, Mountain View, California); the ALIGN program (Version 2.0) and GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Version 8 (available from
10 Genetics Computer Group (GCG), 575 Science Drive, Madison, Wisconsin, USA). Alignments using these programs can be performed using the default parameters. A PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used with the ALIGN program when comparing amino acid sequences. BLAST nucleotide searches can be performed with the BLASTN program, score =
15 100, wordlength = 12, to obtain nucleotide sequences homologous to a nucleotide sequence encoding a protein of the invention. BLAST protein searches can be performed with the BLASTX program, score = 50, wordlength = 3, to obtain amino acid sequences homologous to a protein or polypeptide of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST (in BLAST
20 2.0) can be utilized. Alternatively, PSI-BLAST (in BLAST 2.0) can be used to perform an iterated search that detects distant relationships between molecules. When utilizing BLAST, Gapped BLAST, PSI-BLAST, the default parameters of the respective programs (e.g., BLASTN for nucleotide sequences, BLASTX for proteins) can be used. See <http://www.ncbi.nlm.nih.gov>. Alignment may also be
25 performed manually by inspection. For purposes of the present invention, comparison of nucleotide or protein sequences for determination of percent sequence identity to the sequences described herein is preferably made using the Gapped BLAST program (Version 2.0 or later) with its default parameters or any equivalent program. By "equivalent program" is intended any sequence
30 comparison program that, for any two sequences in question, generates an alignment having identical nucleotide or amino acid residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by the preferred program.

By "functionally equivalent" is intended that the sequence of the variant defines a chain that produces a protein having substantially the same biological effect as the native protein of interest. Such functionally equivalent variants that comprise substantial sequence variations are also encompassed by the invention.

5 Thus, for purposes of the present invention, a functionally equivalent variant of GPP synthase will catalyze the formation of GPP from dimethylallyl diphosphate (DMAPP) and isopentenyl diphosphate (IPP); that of limonene synthase will catalyze the formation of limonene from GPP; that of carveol synthase will catalyze the formation of carveol from limonene; and that of S-linalool synthase

10 will catalyze the formation of S-linalool from GPP.

In view of the ability to transform crop plants to express various heterologous compounds, it would be desirable to transform maize plants to express monoterpene pathway enzymes and thereby produce effective amounts of the monoterpenes limonene, carveol and/or S-linalool so that by consuming the

15 tissues of the plant an insect, such as larvae of Corn Rootworm, would consume insecticidally effective amounts of the monoterpenes; or be caused to avoid feeding on the plant.

While most gene products are peptides, a monoterpene is not a peptide or peptide derivative and is not expressed from genes in the form of a peptide or

20 peptide derivative, but is produced enzymatically as a secondary metabolite within the cells of some plants. It is determined that the biosynthetic apparatus necessary for the production of the monoterpene limonene may not be present in many plant cells which do not produce limonene, or may not produce detectable, insecticidally effective amounts of limonene, and this appears to include maize cells. Such plant

25 cells must be engineered with at least one enzyme which can be produced through the expression of exogenous (heterologous) genes. One such enzyme is limonene synthase, also known as limonene cyclase, which can directly synthesize limonene from geranyl pyrophosphate (GPP), which is found widely in both procaryotic and eucaryotic cells, although, as discussed below, is in some cases not produced

30 in quantities sufficient to make insecticidally effective amounts of limonene.

Since genes which code for a monoterpene synthase can be synthesized, either directly using a DNA sequence obtained by working backwards from the known amino acid sequence of a particular monoterpene synthase and preferably

using plant-preferred codons, or by cloning from natural sources of monoterpenes, the resulting sequence can be inserted into an appropriate expression cassette, and introduced into cells of a susceptible plant species or a suitable endophytic bacterium, so that an especially preferred embodiment of this method involves
5 inserting into the genome of the plant or bacterium a DNA sequence coding for a monoterpene synthase, in proper reading frame relative to transcription initiator and promoter sequences active in the plant or bacterium. Transcription and translation of the coding sequence under control of the regulatory sequences, can cause expression of the enzyme at levels which provide an effective amount of a
10 monoterpene such as limonene in the tissue of the plant which are normally infested by the larvae.

As an illustration, it can be noted that Colby *et al.*, at the Keystone Symposium on Crop Improvement via Biotechnology: An International Perspective, Keystone, Colorado, USA, April 10-16, 1992, as reported in *J. Cell*
15 *Biochem. Suppl.* 16 F, 230 (1992), have isolated and characterized cDNA encoding limonene cyclase from spearmint. To isolate and study the gene(s) (sic) encoding limonene synthase and to produce enough of the enzyme for structural studies, they used standard methods to extract RNA from young leaves of *Mentha spicata* and constructed a cDNA library in λ ZAP XR (Stratagene) from poly (A) + RNA.
20 They designed three degenerate oligonucleotides based on internal amino acid sequences obtained from Edman degradation of purified limonene synthase and screened 250,000 clones to identify six positive clones that hybridized to all three oligonucleotides. The resulting clones could be used in the methods of this invention which involve plant transformation. However, Colby *et al.* indicate no
25 appreciation of the value of the enzyme in conferring resistance to insects in plants.

In certain plants, including maize, at least one additional gene encoding GPP synthase is required for generation of plants with resistance to insects. Due to the fact that natural levels of GPP are low in such plants, there may be inadequate amounts of GPP for limonene production in these plants when they are
30 transformed solely with the limonene synthase gene. In contrast, GPP levels in other plant species, such as spearmint, is adequate for limonene production. In such species, some GPP is used to generate limonene, some to generate other metabolites. Because GPP is derived from a pathway that is common among plant

species, the introduction of the GPP synthase gene and at least one monoterpene synthase gene such as limonene synthase into plant species lacking GPP can generate transgenic plants capable of producing GPP, and a desired monoterpene at levels effective to confer resistance to insects.

5 In this manner, the invention encompasses transforming plants with nucleotide sequences encoding GPP synthase and limonene synthase for the generation of plants producing effective amounts of limonene; with nucleotide sequences encoding GPP synthase, limonene synthase and carveol synthase for the generation of plants producing effective amounts of carveol; and with nucleotide
10 sequences encoding GPP synthase and S-linalool synthase for the generation of plants producing effective amounts of S-linalool.

In one embodiment, the plant which can be benefitted by this invention is preferably a plant susceptible to infestation and damage by the larvae of the genus *Diabrotica* or whose harvested material is subject to attack by larvae of that insect.
15 A prime example is corn (*Zea mays*). Thus the methods of this invention are readily applicable via conventional techniques to numerous plant species, particularly those susceptible to *Diabrotica spp.*, including, without limitation, species from the genera *Fragaria*, *Lotus*, *Medicago*, *Onobrychis*, *Trifolium*, *Trigonella*, *Vigna*, *Citrus*, *Linum*, *Geranium*, *Manicot*, *Daucus*, *Arabidopsis*,
20 *Brassica*, *Raphanus*, *Sinapis*, *Atropa*, *Capsicum*, *Datura*, *Hyoscyamus*, *Lycopersicon*, *Nicotiana*, *Solanum*, *Petunia*, *Digitalis*, *Majorana*, *Cichorium*, *Helianthus*, *Lactuca*, *Bromus*, *Asparagus*, *Antirrhinum*, *Hemerocallis*, *Nemesia*, *Pelargonium*, *Panicum*, *Pennisetum*, *Ranunculus*, *Senecio*, *Salpiglossis*, *Cucumis*, *Browallia*, *Glycine*, *Lolium*, *Triticum*, and *Datura*.

25 Thus, the methods of the invention are useful to transform plants and create or enhance resistance to insects in plants. By resistance to insects is intended that the plant-insect interaction, preferably a plant-insect pest interaction, is reduced, ameliorated, prevented, or eliminated.

Insect pests include but are not limited to insects selected from the orders
30 Coleoptera, Diptera, Hymenoptera, Lepidoptera, Mallophaga, Homoptera, Hemiptera, Orthoptera, Thysanoptera, Dermaptera, Isoptera, Anoplura, Siphonaptera, Trichoptera, etc., particularly Coleoptera and Lepidoptera. Insect pests of the invention for the major crops include: Maize: *Ostrinia nubilalis*,

- European corn borer; *Agrotis ipsilon*, black cutworm; *Helicoverpa zea*, corn earworm; *Spodoptera frugiperda*, fall armyworm; *Diatraea grandiosella*, southwestern corn borer; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Diatraea saccharalis*, sugarcane borer; *Diabrotica virgifera*, western corn rootworm; *Diabrotica longicornis barberi*, northern corn rootworm; *Diabrotica undecimpunctata howardi*, southern corn rootworm; *Melanotus spp.*, wireworms; *Cyclocephala borealis*, northern masked chafer (white grub); *Cyclocephala immaculata*, southern masked chafer (white grub); *Popillia japonica*, Japanese beetle; *Chaetocnema pulicaria*, corn flea beetle; *Sphenophorus maidis*, maize billbug; *Rhopalosiphum maidis*, corn leaf aphid; *Anuraphis maidiradicis*, corn root aphid; *Blissus leucopterus leucopterus*, chinch bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus sanguinipes*, migratory grasshopper; *Hylemya platura*, seedcorn maggot; *Agromyza parvicornis*, corn blot leafminer; *Anaphothrips obscurus*, grass thrips; *Solenopsis milesta*, thief ant; *Tetranychus urticae*, twospotted spider mite; Sorghum: *Chilo partellus*, sorghum borer; *Spodoptera frugiperda*, fall armyworm; *Helicoverpa zea*, corn earworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Feltia subterranea*, granulate cutworm; *Phyllophaga crinita*, white grub; *Eleodes*, *Conoderus*, and *Aeolus spp.*, wireworms; *Oulema melanopus*, cereal leaf beetle; *Chaetocnema pulicaria*, corn flea beetle; *Sphenophorus maidis*, maize billbug; *Rhopalosiphum maidis*, corn leaf aphid; *Sipha flava*, yellow sugarcane aphid; *Blissus leucopterus leucopterus*, chinch bug; *Contarinia sorghicola*, sorghum midge; *Tetranychus cinnabarinus*, carmine spider mite; *Tetranychus urticae*, twospotted spider mite; Wheat: *Pseudaletia unipunctata*, army worm; *Spodoptera frugiperda*, fall armyworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Agrotis orthogonia*, western cutworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Oulema melanopus*, cereal leaf beetle; *Hypera punctata*, clover leaf weevil; *Diabrotica undecimpunctata howardi*, southern corn rootworm; Russian wheat aphid; *Schizaphis graminum*, greenbug; *Macrosiphum avenae*, English grain aphid; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Melanoplus sanguinipes*, migratory grasshopper; *Mayetiola destructor*, Hessian fly; *Sitodiplosis mosellana*, wheat midge; *Meromyza americana*, wheat stem maggot; *Hylemya coarctata*, wheat bulb fly; *Frankliniella*

fusca, tobacco thrips; *Cephus cinctus*, wheat stem sawfly; *Aceria tulipae*, wheat curl mite; Sunflower: *Suleima helianthana*, sunflower bud moth; *Homoeosoma electellum*, sunflower moth; *zygogramma exclamationis*, sunflower beetle; *Bothyrus gibbosus*, carrot beetle; *Neolasioptera murtfeldtiana*, sunflower seed
5 midge; Cotton *Heliothis virescens*, cotton budworm; *Helicoverpa zea*, cotton bollworm; *Spodoptera exigua*, beet armyworm; *Pectinophora gossypiella*, pink bollworm; *Anthonomus grandis*, boll weevil; *Aphis gossypii*, cotton aphid; *Pseudatomoscelis seriatus*, cotton fleahopper; *Trialeurodes abutilonea*, bandedwinged whitefly; *Lygus lineolaris*, tarnished plant bug; *Melanoplus*
10 *femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Thrips tabaci*, onion thrips; *Frankliniella fusca*, tobacco thrips; *Tetranychus cinnabarinus*, carmine spider mite; *Tetranychus urticae*, twospotted spider mite; Rice: *Diatraea saccharalis*, sugarcane borer; *Spodoptera frugiperda*, fall armyworm; *Helicoverpa zea*, corn earworm; *Colaspis brunnea*, grape colaspis;
15 *Lissorhoptrus oryzophilus*, rice water weevil; *Sitophilus oryzae*, rice weevil; *Nephotettix nigropictus*, rice leafhopper; *Blissus leucopterus leucopterus*, chinch bug; *Acrosternum hilare*, green stink bug; Soybean: *Pseudoplusia includens*, soybean looper; *Anticarsia gemmatilis*, velvetbean caterpillar; *Plathypena scabra*, green cloverworm; *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm;
20 *Spodoptera exigua*, beet armyworm; *Heliothis virescens*, cotton budworm; *Helicoverpa zea*, cotton bollworm; *Epilachna varivestis*, Mexican bean beetle; *Myzus persicae*, green peach aphid; *Empoasca fabae*, potato leafhopper; *Acrosternum hilare*, green stink bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Hylemya platura*, seedcorn maggot;
25 *Sericothrips variabilis*, soybean thrips; *Thrips tabaci*, onion thrips; *Tetranychus turkestanii*, strawberry spider mite; *Tetranychus urticae*, twospotted spider mite; Barley: *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Schizaphis graminum*, greenbug; *Blissus leucopterus leucopterus*, chinch bug; *Acrosternum hilare*, green stink bug; *Euschistus servus*,
30 brown stink bug; *Delia platura*, seedcorn maggot; *Mayetiola destructor*, Hessian fly; *Petrobia latens*, brown wheat mite; Oil Seed Rape: *Brevicoryne brassicae*, cabbage aphid; *Phyllotreta cruciferae*, Flea beetle; *Mamestra configurata*, Bertha armyworm; *Plutella xylostella*, Diamond-back moth; *Delia* ssp., Root maggots.

Preferred plants that are to be transformed according to the methods of this invention are cereal crops, including maize, rye, barley, wheat, sorghum, oats, millet, rice, triticale, sunflower, alfalfa, rapeseed and soybean, fiber crops, such as cotton, fruit crops, such as melons, and vegetable crops, including onion, pepper, tomato, cucumber, squash, carrot, crucifer (cabbage, broccoli, cauliflower), eggplant, spinach, potato and lettuce.

While compounds other than monoterpenes have some effective insecticidal activity at high concentrations in pure form, plant cell expression at such high concentrations is either not possible in a living plant cell system, or is not feasible if the commercially useful characteristics of the plant are to be preserved in terms of production of oils, starches, fibers, or other materials. Monoterpenes, on the other hand, are not directly expressed as the gene product, and the peptide or peptides which is or are expressed in the methods of this invention is an enzyme which can catalyze the synthesis of large amounts of a monoterpene(s) in the tissues of the transformed plant (e.g. limonene synthase), and in instances in which it is required, an enzyme which can catalyze the synthesis of large amounts of substrate for a downstream monoterpene synthase (e.g. GPP synthase).

The genes utilized in the invention, including GPP-, carveol, limonene-, and S-linalool synthase genes can be optimized for enhanced expression in plants of interest. See, for example, EPA0359472; WO91/16432; Perlak *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:3324-3328; and Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498. In this manner, the genes can be synthesized utilizing plant-preferred codons. See, for example, Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498, the disclosure of which is incorporated herein by reference. In this manner, synthetic genes can also be made based on the distribution of codons a particular host uses for a particular amino acid. Thus, the nucleotide sequences can be optimized for expression in any plant. It is recognized that all or any part of the gene sequence may be optimized or synthetic. That is, synthetic or partially optimized sequences may also be used.

The methods of the invention encompass utilizing naturally occurring nucleotide sequences encoding GPP synthase, and the monoterpene synthases limonene-, carveol- and S-linalool synthases; or utilizing synthetically derived

sequences encoding these proteins. The naturally occurring nucleotide sequences utilized in the methods of the invention are set forth in Figures 6, 8, 10 and 12.

The methods also encompass utilizing nucleotide sequences isolated from various organisms including plants by hybridization with partial sequences
5 obtained from the natural sequences as set forth above. Conditions that will permit other DNA sequences to hybridize to the DNA sequences set forth herein can be determined in accordance with techniques generally known in the art. For example, hybridization of such sequences may be carried out under conditions of reduced stringency, medium stringency, or high stringency conditions (*e.g.*,
10 conditions represented by a wash stringency of 35-40% Formamide with 5x Denhardt's solution, 0.5% SDS, and 1x SSPE at 37°C; conditions represented by a wash stringency of 40-45% Formamide with 5x Denhardt's solution, 0.5% SDS, and 1x SSPE at 42°C; and conditions represented by a wash stringency of 50% Formamide with 5x Denhardt's solution, 0.5% SDS, and 1x SSPE at 42°C,
15 respectively. See Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York). The methods further encompass utilizing nucleotide sequences isolated from various organisms including plants by other well known methods such as PCR using the natural sequences as set forth above.

20 The methods of the invention comprise utilizing expression cassettes with constitutive or tissue-specific promoters. Promoters that may be used in the expression cassettes include without limitation nos, ocs, phaseolin, FMV and other promoters isolated from the DNA of plants or other sources, both natural and synthetic.

25 Constitutive promoters would provide a constant production of the enzymes GPP-, limonene-, carveol and/or S-linalool synthase and thereby the corresponding monoterpene(s). Such constitutive promoters include, for example, the core promoter of the Rsyn7 (U.S. Patent Application Serial No. 08/661,601), the core CaMV 35S promoter (Odell *et al.* (1985) *Nature* 313:810-812); rice actin
30 (McElroy *et al.* (1990) *Plant Cell* 2:163-171); ubiquitin (Christensen *et al.* (1989) *Plant Mol. Biol.* 12:619-632 and Christensen *et al.* (1992) *Plant Mol. Biol.* 18 675-689); pEMU (Last *et al.* (1991) *Theor. Appl. Genet.* 81:581-588); MAS (Velten *et al.* (1984) *EMBO J.* 3:2723-2730); ALS promoter (U.S. Patent Application Serial

No. 08/409,297), and the like. Other constitutive promoters include, for example, U.S. Patent Nos. 5,608,149; 5,608,144; 5,604,121; 5,569,597; 5,466,785; 5,399,680; 5,268,463; and 5,608,142.

5 An efficient plant promoter that may be used is an overproducing plant promoter. Overproducing plant promoters that may be used in this invention include the promoter of the small sub-unit (ss) of the ribulose-1,5-bisphosphate carboxylase from soybean (Berry-Lowe *et al.* (1982) *J. Mol. and App. Gen.* 1:483-498, and the promoter of the chlorophyll a-b binding protein. However, these two promoters are known to be light-induced in eukaryotic plant cells (see, for
10 example, *Genetic Engineering of Plants, An Agricultural Perspective*, Cashmore, Pelham, New York, 1983, pp. 29-38, G. Coruzzi *et al.* (1983) *J. Biol. Chem.* 258:1399 and P. Dunsmuir *et al.* (1983) *J. Mol. and App. Gen.* 2:285 and may be less desirable when root expression is desired.

A tissue-specific promoter (or promoters) can be used in any instance
15 where it may be desirable to localize production of the desired monoterpene to an insect-infested tissue or to a tissue which is efficient in production of a desired enzyme. The utilization of tissue-specific promoters would increase or decrease the expression of monoterpene synthases and production of the corresponding monoterpenes in specific tissues of the plant. It is recognized that in manipulating
20 the level of monoterpene production as such, it may be desirable to increase or decrease the levels of such molecules in a particular tissue, since tolerance of various tissues to increased expression of specific monoterpenes may vary. Thus, it may be desirable to increase expression in selected tissues, or at varying levels in different tissues by the use of tissue-specific promoters.

25 Particular tissue-specific promoters of interest includes root-preferred promoters. The utilization of such promoters would provide a method of selectively creating or enhancing resistance to insects, and/or manipulating levels of monoterpene molecules in the root. Since corn rootworm attack roots, root-specific promoters are especially preferred for the control of corn rootworm, while
30 minimizing limonene production in the agronomically valuable parts of the plant. Such selectivity could also be particularly desirable in plants in which the root constitutes the food crop, including, but not limited to carrot, potato, radish, and the like.

Root specific promoters are known and can be selected from the many available from the literature or isolated de novo from various compatible species. For example, Hire, *et al.* (1992) *Plant Mol Biology*, 20(2): 207-218, describe a root-specific glutamine synthetase gene from soybean. Keller and Baumgartner, (1991) *The Plant Cell*, 3(10):1051-1061, describe a root-specific control element in the GRP 1.8 gene of French bean. Sanger *et al.* (1990) *Plant Mol. Biology*, 14(3): 433-443, discuss the root-specific promoter of the Mannopine Synthase (MAS) gene of *Agrobacterium tumefaciens*. Miao *et al.* (1991) *The Plant Cell*, 3(1):11-22, describe a full-length cDNA clone encoding cytosolic glutamine synthetase (GS), which is expressed in roots and root nodules of soybean. Bogusz *et al.* (1990) *The Plant Cell*, 2(7):633-641, discusses two root-specific promoters isolated from hemoglobin genes from the nitrogen-fixing nonlegume *Parasponia andersonii* and the related non-nitrogen-fixing nonlegume *Trema tomentosa*. The promoters of these genes were linked to a β -glucuronidase reporter gene and introduced into both the nonlegume *Nicotiana tabacum* and the legume *Lotus corniculatus*, and in both instances root-specific promoter activity was preserved. Leach and Aoyagi (1991) *Plant Science (Limerick)* 79(1):69-76, describe their analysis of the promoters of the highly expressed rolC and rolD root-inducing genes of *Agrobacterium rhizogenes*. They concluded that enhancer and tissue-specific DNA determinants are dissociated in those promoters. Teeri *et al.* (1989) *EMBO Journal*, 8(2):343-350, used gene fusion to lacZ to show that the *Agrobacterium* T-DNA gene encoding octopine synthase is especially active in the epidermis of the root tip and that the TR2' gene was root specific in the intact plant and stimulated by wounding in leaf tissue, an especially desirable combination of characteristics for use with an insecticidal or larvicidal gene. The TR1' gene, fused to NPTII, (neomycin phosphotransferase II) showed similar characteristics. Additional root-preferred promoters include the VfENOD-GRP3 gene promoter (Kuster H *et al.* (1995) *Plant Mol. Biol.* 29(4):759-772); and rolB promoter (Capana *et al.* (1994) *Plant Mol. Biol.* 25(4):681-691. See also U.S. Patent Nos. 5,633,363; 5,459,252; 5,401,836; 5,110,732; and 5,023,179.

Other tissue-specific promoters include those described in Yamamoto *et al.* (1997) *Plant J.* 12(2):255-265; Kawamata *et al.* (1997) *Plant Cell Physiol.* 38(7):792-803; Hansen *et al.* (1997) *Mol. Gen. Genet.* 254(3):337-343; Russell *et*

al. (1997) *Transgenic Res.* 6(2):157-168; Rinehart *et al.* (1996) *Plant Physiol.* 112(3):1331-1341; Van Camp *et al.* (1996) *Plant Physiol.* 112(2):525-535; Canevascini *et al.* (1996) *Plant Physiol.* 112(2):513-524; Yamamoto *et al.* (1994) *Plant Cell Physiol.* 35(5):773-778; Lam (1994) *Results Probl. Cell Differ.* 20:181-196; Orozco *et al.* (1993) *Plant Mol. Biol.* 23(6):1129-1138; Matsuoka *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90(20):9586-9590; and Guevara-Garcia *et al.* (1993) *Plant J.* 4(3):495-505.

Seed-preferred promoters includes both seed-specific promoters (those promoters active during seed development) as well as seed-germinating promoters (those promoters active during seed germination). Such promoters include Cim1 (cytokinin-induced message); cZ19B1 (maize 19KDa zein); milps (myo-inositol-1-phosphate synthase); celA (cellulose synthase); end1 (*Hordeum vulgare* mRNA clone END1); and imp3 (myo-inositol monophosphate-3). For dicots, particular promoters include phaseolin, napin, β -conglycinin, soybean lectin, and the like. For monocots, particular promoters include maize 15Kd zein, 22KD zein, 27kD zein, waxy, shrunken 1, shrunken 2, globulin 1, etc.

In native tissues which synthesize monoterpenes, GPP-, limonene- and S-linalool synthases are targeted to plastids via plastid targeting sequences (transit peptides) typically contained at the N-terminus of the preprotein forms of the enzymes. The targeting sequences are then cleaved to release the mature enzymes in plastids. Such transit peptides can be identified in the primary amino acid sequences of the preproteins by those ordinarily skilled in the art. For example, see Colby *et al.* (1993) *J. Biol. Chem.* 268(31):23016-23024, for the transit peptide sequence of limonene synthase. Native carveol synthase is associated with the endoplasmic reticulum, while native carveol dehydrogenase could be cytoplasmic.

Furthermore, the substrates of GPP synthase, IPP (IDP) and DMAPP (DMADP) can be found in plastids or in the cytoplasm (Bohlmann *et al.* (1998) *Proc. Natl. Acad. Sci. USA* 95:4126-4133). In plastids, these substrates are most likely synthesized via the glyceraldehyde-3-phosphate/pyruvate (Rohmer) pathway, and in the cytosol via the mevalonate pathway in the cytosol or the mevalonate pathway.

Thus, for manipulating a metabolic pathway involving monoterpene production, it would be beneficial to target GPP-, limonene-, and S-linalool

synthase to plastids; and carveol synthase and dehydrogenase to the endoplasmic reticulum (ER). This targeting could be achieved by use of the native targeting sequences contained in the sequences of the native proteins, or by addition or exchange of heterologous subcellular targeting signals. Alternatively, the enzymes
5 utilized in the methods of the invention could be directed to the cytoplasm by deletion of the plastid and/or ER targeting signals. Methods for deletion, exchange and addition of nucleotide sequences are well known in the art, and can be readily used for manipulation of nucleotide segments encoding targeting signals of interest as described herein.

10 Heterologous sequences which can be used to target the desired enzymes of the invention to plastids include chloroplast targeting sequences. Chloroplast targeting sequences are known in the art and include the chloroplast small subunit of ribulose-1,5-bisphosphate carboxylase (Rubisco), (de Castro Silva Filho *et al.* (1996) *Plant Mol. Biol.* 30:769-780; Schnell, *et al.* (1991) *J. Biol. Chem.* 266(5):3335-3342), 5-(enolpyruvyl)shikimate-3-phosphate synthase (EPSPS)
15 (Archer *et al.* (1990) *J. Bioenerg. Biomemb.* 22(6):789-810); tryptophan synthase (Zhao *et al.* (1995) *J. Biol. Chem.* 270(11):6081-6087); plastocyanin (Lawrence *et al.* (1997) *J. Biol. Chem.* 272(33):20357-20363); chorismate synthase (Schmidt *et al.* (1993) *J. Biol. Chem.* 268(36):27477-27457); and the light harvesting
20 chlorophyll a/b binding protein (LHBP) (Lamppa *et al.* (1988) *J. Biol. Chem.* 263:14996-14999). See also Von Heijne *et al.* (1991) *Plant Mol. Biol. Rep.* 9:104-126; Clark *et al.* (1989) *J. Biol. Chem.* 264:17544-17550; della-Cioppa *et al.* (1987) *Plant Physiol.* 84:965-968, Romer *et al.* (1993) *Biochem. Biophys. Res Commun.* 196:1414-1421; and Shah *et al.* (1986) *Science* 233:478-481.

25 Likewise, methods for transformation of chloroplasts are known in the art. See, for example, Svab *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:8526-8530; Svab and Maliga (1993) *Proc. Natl. Acad. Sci. USA* 90:913-917; Staub and Maliga (1993) *Embo J.* 12:601-606. The method relies on particle gun delivery of DNA containing a selectable marker and targeting of the DNA to the plastid genome
30 through homologous recombination. Additionally, plastid transformation can be accomplished by transactivation of a silent plastid-borne transgene by tissue-specific expression of a nuclear-encoded and plastid-directed RNA polymerase.

Such a system has been reported in McBride *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:7301-7305.

The nucleotide sequences utilized in methods of the invention can be introduced into any plant. The sequences to be introduced may be used in
5 expression cassettes for expression in any plant of interest where expression in the plant is necessary for transcription.

While it may be preferable to express the sequences using heterologous promoters, the native promoter sequences may be used. Such constructs would change expression levels of the enzymes of the invention in the plant or plant cell.
10 Thus, the phenotype of the plant or plant cell is altered.

Where expression cassettes are needed, such expression cassettes will comprise a transcriptional initiation region linked to the coding sequence or antisense sequence of the nucleotide of interest. Such an expression cassette is provided with a plurality of restriction sites for insertion of the sequence to be
15 under the transcriptional regulation of the regulatory regions. The expression cassette may additionally contain selectable marker genes.

The transcriptional initiation region, the promoter, may be native or analogous or foreign or heterologous to the plant host. Additionally, the promoter may be the natural sequence or alternatively a synthetic sequence. By foreign is
20 intended that the transcriptional initiation region is not found in the native plant into which the transcriptional initiation region is introduced. As used herein a chimeric gene comprises a coding sequence operably linked to a transcription initiation region that is heterologous to the coding sequence.

The transcriptional cassette will include in the 5'-to-3' direction of
25 transcription, a transcriptional and translational initiation region, a DNA sequence of interest, and a transcriptional and translational termination region functional in plants. The termination region may be native with the transcriptional initiation region, may be native with the DNA sequence of interest, or may be derived from another source. Convenient termination regions are available from the Ti-plasmid
30 of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also Guerineau *et al.* (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon *et al.* (1991) *Genes Dev.* 5:141-149; Mogen *et al.* (1990) *Plant Cell.* 2:1261-1272; Munroe *et al.* (1990) *Gene* 91:151-

158; Ballas *et al.* (1989) *Nucleic Acids Res.* 17:7891-7903; Joshi *et al.* (1987) *Nucleic Acids Res.* 15:9627-9639.

The methods of the invention encompass use of expression cassettes for expression of nucleotide sequences encoding GPP- and monoterpene synthases in
5 the plant of interest. The cassette will include 5' and 3' regulatory sequences operably linked to the sequence of interest. The cassette may additionally contain at least one additional sequence to be cotransformed into the organism. Alternatively, the additional sequence(s) can be provided on another expression cassette.

10 Where appropriate, sequences utilized in the methods of the invention and additional gene(s) may be optimized for increased expression in the transformed plant. That is, these nucleotide sequences can be synthesized using plant preferred codons for improved expression. Methods are available in the art for synthesizing plant preferred genes. See, for example, U.S. Patent Nos. 5,380,831, 5,436, 391,
15 and Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498, herein incorporated by reference.

Additional sequence modifications are known to enhance gene expression in a cellular host. These include elimination of sequences encoding spurious polyadenylation signals, exon-intron splice site signals, transposon-like repeats,
20 and other such well-characterized sequences which may be deleterious to gene expression. The G-C content of the sequence may be adjusted to levels average for a given cellular host, as calculated by reference to known genes expressed in the host cell. When possible, the sequence is modified to avoid predicted hairpin secondary mRNA structures.

25 The expression cassettes may additionally contain 5' leader sequences in the expression cassette construct. Such leader sequences can act to enhance translation. Translation leaders are known in the art and include: picornavirus leaders, for example, EMCV leader (Encephalomyocarditis 5' noncoding region) (Elroy-Stein *et al.* (1989) *Proc. Nat. Acad. Sci. USA* 86:6126-6130); potyvirus
30 leaders, for example, TEV leader (Tobacco Etch Virus) (Allison *et al.* (1986)); MDMV leader (Maize Dwarf Mosaic Virus) (*Virology* 154:9-20); human immunoglobulin heavy-chain binding protein (BiP) (Macejak and Sarnow (1991) *Nature* 353:90-94); untranslated leader from the coat protein mRNA of alfalfa

mosaic virus (AMV RNA 4) (Jobling and Gehrke (1987) *Nature* 325:622-625); tobacco mosaic virus leader (TMV) (Gallie *et al.* (1989) *Molecular Biology of RNA*, pages 237-256); and maize chlorotic mottle virus leader (MCMV) (Lommel *et al.* (1991) *Virology* 81:382-385). See also Della-Cioppa *et al.* (1987) *Plant Physiology* 84:965-968. Other methods known to enhance translation can also be utilized, for example, introns, and the like.

In preparing the expression cassette, the various DNA fragments may be manipulated, so as to provide for the DNA sequences in the proper orientation and, as appropriate, in the proper reading frame. Toward this end, adapters or linkers may be employed to join the DNA fragments or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous DNA, removal of restriction sites, or the like. For this purpose, *in vitro* mutagenesis, primer repair, restriction, annealing, resubstitutions, *e.g.*, transitions and transversions, may be involved.

The methods of the invention can be used in transforming or transfecting any plant. In this manner, genetically modified plants, plant cells, plant tissue, seed, and the like can be obtained. Transformation protocols as well as protocols for introducing nucleotide sequences into plants may vary depending on the type of plant or plant cell, *i.e.* monocot or dicot, targeted for transformation. Suitable methods of introducing nucleotide sequences into plant cells and subsequent insertion into the plant genome include microinjection (Crossway *et al.* (1986) *Biotechniques* 4:320-334), electroporation (Riggs *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606, *Agrobacterium*-mediated transformation (Hinchee *et al.* (1988) *Biotechnology* 6:915-921), direct gene transfer (Paszkowski *et al.* (1984) *EMBO J.* 3:2717-2722), and ballistic particle acceleration (see, for example, Sanford *et al.*, U.S. Patent No. 4,945,050; Tomes *et al.* (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips (Springer-Verlag, Berlin); and McCabe *et al.* (1988) *Biotechnology* 6:923-926). Also see Weissinger *et al.* (1988) *Annual Rev. Genet.* 22:421-477; Sanford *et al.* (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou *et al.* (1988) *Plant Physiol.* 87:671-674 (soybean); McCabe *et al.* (1988) *Bio/Technology* 6:923-926 (soybean); Finer and McMullen (1991) *In Vitro Cell Dev. Biol.*

27P:175-182 (soybean); Singh *et al.* (1998) *Theor. Appl. Genet.* 96:319-324 (soybean); Datta *et al.* (1990) *Biotechnology* 8:736-740 (rice); Klein *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-4309 (maize); Klein *et al.* (1988) *Biotechnology* 6:559-563 (maize); Tomes *et al.* "Direct DNA Transfer into Intact
5 Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg (Springer-Verlag, Berlin) (1995) (maize); Klein *et al.* (1988) *Plant Physiol.* 91:440-444 (maize); Fromm *et al.* (1990) *Biotechnology* 8:833-839 (maize); Hooydaas-Van Slogteren and Hooykaas (1984) *Nature (London)* 311:763-764; Bytebier *et al.* (1987) *Proc. Natl. Acad. Sci.*
10 *USA* 84:5345-5349 (Liliaceae); De Wet *et al.* (1985) in *The Experimental Manipulation of Ovule Tissues*, ed. Chapman *et al.* (Longman, New York), pp. 197-209 (pollen); Kaeppler *et al.* (1990) *Plant Cell Reports* 9:415-418; and Kaeppler *et al.* (1992) *Theor. Appl. Genet.* 84:560-566 (whisker-mediated transformation); D'Halluin *et al.* (1992) *Plant Cell* 4:1495-1505 (electroporation);
15 Li *et al.* (1993) *Plant Cell Reports* 12:250-255 and Christou and Ford (1995) *Annals of Botany* 75:407-413 (rice); Osjoda *et al.* (1996) *Nature Biotechnology* 14:745-750 (maize via *Agrobacterium tumefaciens*); all of which are herein incorporated by reference. Thus, the invention provides a method for imparting insect resistance in *Agrobacterium tumefaciens*-susceptible plants in which the
20 expression cassette is introduced into the cells by infecting the cells with *Agrobacterium tumefaciens*, a plasmid of which has been modified to include a plant expression cassette which expresses GPP- and/or a monoterpene synthase in the manner of this invention.

The modified plant may be grown into plants in accordance with
25 conventional ways. See, for example, McCormick *et al.* (1986) *Plant Cell. Reports* 5:81-84. These plants may then be grown, and either pollinated with the same transformed strain or different strains, and the resulting plants or hybrid plants the desired phenotypic characteristic identified. Two or more generations may be grown to ensure that the subject phenotypic characteristic is stably maintained and
30 inherited and then seeds harvested to ensure the desired phenotype or other property has been achieved.

In a preferred embodiment, the invention provides methods of imparting resistance to *Diabrotica spp.* to plants of a susceptible taxon, comprising the steps of:

- (a) culturing cells or tissues from at least one plant from the taxon,
- 5 (b) introducing into the cells of the cell or tissue culture at least one copy of an expression cassette comprising a structural gene coding for at least one monoterpene synthase or both a monoterpene synthase and GPP synthase, operably linked to a promoter that drives expression in a plant cell, and
- (c) regenerating whole plants from the cell or tissue culture with
- 10 resistance to insects. Once whole plants have been obtained, they can be sexually or clonally reproduced in such manner that at least one copy of the sequence provided by the expression cassette is present in the cells of progeny of the reproduction.

Alternatively, once a single transformed plant has been obtained by the
15 foregoing recombinant DNA method, conventional plant breeding methods can be used to transfer the structural gene or genes and associated regulatory sequences via crossing and backcrossing. Such intermediate methods will comprise the further steps of:

- (a) sexually crossing the plant having resistance to insects with a plant
- 20 from the taxon susceptible to insects;
- (b) recovering reproductive material from the progeny of the cross; and
- (c) growing plants having resistance to insects from the reproductive material. Where desirable or necessary, the agronomic characteristics of the susceptible taxon can be substantially preserved by expanding this method to
- 25 include the further steps of repetitively:
 - (a) backcrossing the progeny having resistance to insects with plants from taxon susceptible to insects; and
 - (b) selecting for the expression of resistance to insects (or an associated marker gene) or monoterpene production among the progeny of the backcross,
 - 30 until the desired percentage of the characteristics of the susceptible taxon are present in the progeny along with the gene imparting resistance to insects.

Insect pests of harvested material, including those of stored grain, can also be targets for the methods of this invention. In view of this, the invention also

provides methods for creating or enhancing resistance to insect pests in harvested materials and products obtained from harvested materials, by expressing at least one monoterpene synthase and/or GPP synthase in the plant such that effective amounts of the desired monoterpene is produced in the harvested material and products obtained from such material.

The following examples are offered by way of illustration and not by way of limitation. The following description further exemplifies the compositions of this invention and the methods of making and using them. However, it will be understood that other methods, known by those of ordinary skill in the art to be equivalent, can also be employed.

EXAMPLES 1-6

Insect larvae inhibition and toxicity assays for

Southern Corn Rootworm (SCR) and European Corn Borer (ECB)

Bioassay diets were prepared as described in Czapla and Lang in "Effect of Plant Lectins on the Larval Development of European Corn Borer (Lepidoptera: Pyralidae) and Southern Corn Rootworm (Coleoptera: Chrysomelidae)", *J. Econ. Entomol.* 83:2480-85 (1990), except that low melting temperature agarose replaced the regular agarose so that the diets could be chilled to 37°C prior to the addition of limonene (one assay used the regular agarose diet).

Results were as follows. In Examples 1-4, the test larvae were Southern Corn Rootworm. In Examples 5-6, the test larvae were European Corn Borer. The results of each experiment represent the average from 16-32 insects. All limonene concentrations (ppm) are by weight. The SCR data indicate that limonene is effective against the larvae, but when limonene was used in the same protocol against ECB, little or no effect was seen.

$$\% \text{ Corrected Mortality} = 100 \times \frac{(\text{mortality of treated} - \text{control})}{(100 - \text{control})}$$

$$\% \text{ Wt. Reduction} = 100 \times \frac{(\text{control weight} - \text{treated weight})}{(\text{control wt})}$$

Example:	<u>1</u>		<u>2</u>	
Limonene	% Corr.	% weight	% Corr.	% weight
<u>ppm</u>	<u>Mortal.</u>	<u>Reduction</u>	<u>Mortal.</u>	<u>Reduction</u>
10,000	57	64	80	86
1,000	49	52	19	30
100	57	52	15	6

Example:	<u>3</u>		<u>4</u>	
Limonene	% Corr.	% weight	% Corr.	% weight
<u>ppm</u>	<u>Mortal.</u>	<u>Reduction</u>	<u>Mortal.</u>	<u>Reduction</u>
10,000	96	64	44	0
1,000	7	0	22	0
100	26	0	19	0

Average SCR Results:

Limonene	% Corr.	% weight
<u>ppm</u>	<u>Mortal.</u>	<u>Reduction</u>
10,000	69	54
1,000	24	21
100	29	15

5

Example:	<u>5</u>		<u>6</u>	
Limonene	% Corr.	% weight	% Corr.	% weight
<u>ppm</u>	<u>Mortal.</u>	<u>Reduction</u>	<u>Mortal.</u>	<u>Reduction</u>
10,000	0	28	0	0
1,000	0	16	0	0
100	0	1	0	7

Average ECB Results:

Limonene	% Corr.	% weight
<u>ppm</u>	<u>Mortal.</u>	<u>Reduction</u>
10,000	0	14
1,000	0	8
100	0	4

EXAMPLE 7

Maize callus cultures were transformed by microprojectile bombardment using plasmids containing a cloned gene coding for the limonene synthase (limonene cyclase) enzyme driven by a ubiquitin promoter and a ubiquitin intron and followed downstream by a PIN-II terminator. Whole, fertile plants were regenerated from the transformed callus and analyzed for limonene synthase and limonene. Representative results from one series were as follows:

10

<u>Clone #</u>	<u>Callus LS</u> ¹	<u>Callus</u> <u>Limonene</u> ²	<u>Plant</u>	<u>Plant LS</u>	<u>Plant</u> <u>Limonene</u>
C6	22539	ND	3	1950	ND
			7	2000	ND
C19	9900	ND	6	2100	ND
C3	2550	ND	3	1900	ND
			4	1650	ND

In other experiments, transgenic maize plants, and tissues that exhibited high expression of the limonene synthase protein were produced. The enzyme was extracted from transformed plants and tissues and allowed to react with tritium-labeled geranyl pyrophosphate (GPP). The extracted enzyme converted GPP to limonene, showing that a functional enzyme was being produced. In addition, western blots were done and confirmed the presence of LS protein in transgenic

1 LS = Limonene Synthase

2 ND = not detected above 0.5 ng/g fresh wt.

tissues but not in negative control tissues. Seed was collected from the transformed plants. In sum, whole fertile, transformed plants have been produced that express active limonene cyclase (synthase) according to the teaching of the above-identified application.

5 Western blots were also performed on chloroplasts isolated from plants grown from T1 and T2 seed which expressed the mint limonene synthase gene (Figure 6) in leaf tissue. Thus, the mint limonene synthase was properly targeted to maize plastids and the protein was processed to the correct size, indicating that the mint limonene synthase plastid targeting sequence can be used to target
10 proteins to maize plastids.

EXAMPLE 8

In view of the results set forth in Example 7, demonstrating expression of active limonene synthase in whole, fertile, transformed plants, larvicidally
15 effective amounts of limonene can be produced in such transgenic plants where sufficient substrate is present for the limonene synthase enzyme to act on. Accordingly, maize callus cultures are transformed by art recognized microprojectile bombardment methods using plasmids containing genes coding for both the limonene synthase enzyme and the GPP synthase enzyme, driven, for
20 example, by one or more promoters (a ubiquitin promoter, for example) and followed downstream by, for example, a PIN-II terminator. Whole, fertile plants are regenerated from the transformed callus and analyzed for presence of and/or activity of both enzymes, and are also analyzed for the presence of limonene.

Alternatively, callus is generated from transgenic plants that contain and
25 express the limonene synthase transgene, and such callus cultures are transformed as described above, except that such callus is transformed using plasmids containing a gene coding for the GPP synthase protein. The whole, fertile, transgenic plants regenerated from such transformed callus produce larvicidally effective amounts of limonene.

30

EXAMPLE 9

Average minimum levels of various monoterpenes required for 100% mortality of Western corn rootworm were determined by the methods described in Examples 1-6 as follows:

- 5 Limonene-----200 ppm
 S-linalool---50 ppm
 carveol-----25 ppm

EXAMPLE 10: Construction of Expression Vectors

- 10 Nucleotide sequences encoding limonene, GPP-, carveol and S-linalool synthase, as set forth in SEQ ID NOs: 1, 3, 5, and 7 respectively, are cloned into a plasmid vector, such as that shown in Figure 2, in the sense orientation so that they are under the transcriptional control of the ubiquitin promoter. A selectable marker gene may reside on this plasmid or may be introduced as part of a second plasmid.
- 15 The transformation construct is then available for introduction into maize embryos by bombardment methods as described in Example 12.

20 **EXAMPLE 11: Construction of Expression Vectors**

- Nucleotide sequences encoding GPP- and S-linalool synthase as described in EXAMPLE 10 were cloned in the sense orientation into an expression vector as shown in in Figure 3(PHP14099); such that both coding sequences are under the transcriptional control of the ubiquitin promoter. The plasmid contains the
- 25 selectable marker gene PAT.

- Nucleotide sequences encoding GPP-, limonene- and carveol synthases as described in EXAMPLE 10 were cloned in the sense orientation into an expression vector as shown in Figure 4(PHP14100); such that all three coding sequences are under the transcriptional control of the ubiquitin promoter. The plasmid contains
- 30 the selectable marker gene PAT.

 The transformation constructs PHP14099 and PHP14100 are available for introduction into maize embryos by bombardment methods as described in Example 12.

EXAMPLE 12: Transformation and Regeneration of Maize Callus

Immature maize embryos from greenhouse donor plants are bombarded with a plasmid containing the disclosed GPP- or monoterpene synthase gene operably linked to the ubiquitin promoter plus a plasmid containing the selectable marker gene PAT (Wohlleben et al. (1988) Gene 70:25-37) that confers resistance to the herbicide Bialophos. Transformation is performed as follows. All media recipes are in the Appendix.

10 Preparation of Target Tissue

The ears are surface sterilized in 30% Chlorox bleach plus 0.5% Micro detergent for 20 minutes, and rinsed two times with sterile water. The immature embryos are excised and placed embryo axis side down (scutellum side up), 25 embryos per plate. These are cultured on 560 L medium for 4 days prior to bombardment, in the dark. The day of bombardment, the embryos are transferred to 560 Y medium for 4 hours, arranged within the 2.5-cm target zone.

Preparation of DNA

A plasmid vector comprising the disclosed GPP- or monoterpene synthase operably linked to the ubiquitin promoter is constructed. This plasmid DNA plus plasmid DNA containing a PAT selectable marker is precipitated onto 1.1 μm (average diameter) tungsten pellets using a CaCl_2 precipitation procedure as follows:

- 100 μl prepared tungsten particles in water
- 25 10 μl (1 μg) DNA in TrisEDTA buffer (1 μg total)
- 100 μl 2.5 M CaCl_2
- 10 μl 0.1 M spermidine

Each reagent is added sequentially to the tungsten particle suspension, while maintained on the multitube vortexer. The final mixture is sonicated briefly and allowed to incubate under constant vortexing for 10 minutes. After the precipitation period, the tubes are centrifuged briefly, liquid removed, washed with 500 ml 100% ethanol, and centrifuged for 30 seconds. Again the liquid is

removed, and 105 μ l 100% ethanol is added to the final tungsten particle pellet. For particle gun bombardment, the tungsten/DNA particles are briefly sonicated and 10 μ l spotted onto the center of each macrocarrier and allowed to dry about 2 minutes before bombardment.

5

Particle Gun Treatment

The sample plates are bombarded at level #4 in particle gun #HE34-1 or #HE34-2. All samples receive a single shot at 650 PSI, with a total of ten aliquots taken from each tube of prepared particles/DNA.

10

Subsequent Treatment

Following bombardment, the embryos are kept on 560Y medium for 2 days, then transferred to 560R selection medium containing 3 mg/liter Bialophos, and subcultured every 2 weeks. After approximately 10 weeks of selection, selection-resistant callus clones are sampled for PCR and activity of the acetyl-CoA synthetase gene of interest. Positive lines are transferred to 288J medium to initiate plant regeneration. Following somatic embryo maturation (2-4 weeks), well-developed somatic embryos are transferred to medium for germination and transferred to the lighted culture room. Approximately 7-10 days later, developing plantlets are transferred to medium in tubes for 7-10 days until plantlets are well established. Plants are then transferred to inserts in flats (equivalent to 2.5" pot) containing potting soil and grown for 1 week in a growth chamber, subsequently grown an additional 1-2 weeks in the greenhouse, then transferred to classic 600 pots (1.6 gallon) and grown to maturity. Plants are monitored for expression of the GPP-synthase or monoterpene synthase gene of interest.

25

APPENDIX

Ingredient	Amount	Unit
D-I H ₂ O	900.000	ml
CHU (N6) Basal Salts (SIGMA C-1416)	1.600	g
N6 Macronutrients 10X Stock ##	60.000	ml
Potassium Nitrate	1.680	g
B5H Minor Salts 1000X ###	0.600	ml
B5H Fe Na EDTA 100X ####	6.000	ml
Eriksson's Vitamin Mix (1000X SIGMA-1511)	0.400	ml
S & H Vitamin Mixture 100X Stock (S3766)	6.000	ml
Thiamine.HCL 0.4mg/ml	0.500	ml
L-Proline	1.980	g
Casein Hydrolystate (acid)	0.300	g
Sucrose	20.000	g
Glucose	0.600	g
2,4-D 0.5 mg/ml	1.600	ml
Gelrite @	2.000	g
Dicamba 1 mg/ml #	1.200	ml
Silver Nitrate 2 mg/ml #	1.700	ml

Directions:

@= Add after bringing up to volume

5 #= Add after sterilizing and cooling to temp.

Dissolve ingredients in polished D-I H₂O in sequence

Adjust to pH 5.8

Bring up to volume with polished D-I H₂O after adjusting pH

Sterilize and cool to 60°C.

10 ##= Dissolve 1.660 g of Calcium Chloride Dihydrate in 950.000 ml of polished D-I H₂O. Then dissolve 4.629 of Ammonium Sulfate; 4.000 g of Potassium Phosphate Monobasic KH₂PO₄; 1.850 g of Magnesium Sulfate 7-H₂O, MgSO₄, 7H₂O; and 28.300 g of Potassium Nitrate into sequence. Bring up to volume with polished D-I H₂O.

= Dissolve 3.000 g of Boric Acid; 10.000 g of Manganous Sulfate Monohydrate; 0.250 g of Sodium Molybdate Dihydrate; and 0.750 g of Potassium Iodide in polished D-I H₂O in sequence. Bring up to volume with polished D-I H₂O.

- 5 ##### = Dissolve 3.700 g of Disodium EDTA Dihydrate and 2.790 g of Ferrous Sulfate 7-Hydrate into D-I H₂O. Bring up to volume with D-I H₂O.

Total Volume (L) = 1.00

604 A

Ingredient	Amount	Unit
D-I H ₂ O	900.000	ml
CHU (N6) Basal Salts (SIGMA C-1416)	1.600	g
N6 Macronutrients 10X Stock ##	60.000	ml
Potassium Nitrate	1.680	g
B5H Minor Salts 1000X ###	0.600	ml
B5H Fe Na EDTA 100X ####	6.000	ml
Eriksson's Vitamin Mix (1000X SIGMA-1511)	0.400	ml
S & H Vitamin Mixture 100X Stock (S3766)	6.000	ml
Thiamine.HCL 0.4mg/ml	0.500	ml
L-Proline	1.980	g
Casein Hydrolysate (acid)	0.300	g
Sucrose	20.000	g
Glucose	0.600	g
2, 4-D 0.5mg/ml	1.600	ml
Gelrite @	2.000	g
Dicamba 1mg/ml #	1.200	ml
Silver Nitrate 2mg/ml #	1.700	ml
Bialaphos 1 mg/ml #	3.000	ml

Directions:

@ = Add after bringing up to volume

5 # = Add after sterilizing and cooling to temp.

Dissolve ingredients in polished D-I H₂O in sequence

Adjust to pH 5.8

Bring up to volume with polished D-I H₂O after adjusting pH

Sterilize and cool to 60°C.

10 #### = Dissolve 1.660 g of Calcium Chloride Dihydrate in 950.000 ml of polished D-I H₂O. Then dissolve 4.629 of Ammonium Sulfate, 4.000 g of Potassium Phosphate Monobasic KH₂PO₄, 1.850 g of Magnesium Sulfate 7-H₂O, MgSO₄,

7H₂O; and 28.300 g of Potassium Nitrate into sequence. Bring up to volume with polished D-I H₂O.

- ### = Dissolve 3.000 g of Boric Acid; 10.000 g of Manganous Sulfate Monohydrate; 0.250 g of Sodium Molybdate Dihydrate; and 0.750 g of Potassium Iodide in polished D-I H₂O in sequence. Bring up to volume with polished D-I H₂O.

= Dissolve 3.700 g of Disodium EDTA Dihydrate and 2.790 g of Ferrous Sulfate 7-Hydrate into D-I H₂O. Bring up to volume with D-I H₂O.

Total Volume (L) = 1.00

605 J

Ingredient	Amount	Unit
D-I H ₂ O	900.000	ml
CHU (N6) Basal Salts (SIGMA C-1416)	1.600	g
N6 Macronutrients 10X Stock ##	60.000	ml
Potassium Nitrate	1.680	g
B5H Minor Salts 1000X ###	0.600	ml
B5H Fe Na EDTA 100X ####	6 000	ml
Eriksson's Vitamin Mix (1000X SIGMA-1511)	0 400	ml
S & H Vitamin Mixture 100X Stock (S3766)	6 000	ml
Thiamine.HCL 0.4mg/ml	0 500	ml
Sucrose	20.000	g
Glucose	0 600	g
2, 4-D 0.5mg/ml	1 600	ml
Gelrite @	2 000	g
Dicamba 1mg/ml #	1 200	ml
Silver Nitrate 2mg/ml #	0 425	ml
Bialaphos 1 mg/ml #	3 000	ml

Directions:

@ = Add after bringing up to volume

5 # = Add after sterilizing and cooling to temp.

Dissolve ingredients in polished D-I H₂O in sequence

Adjust to pH 5.8

Bring up to volume with polished D-I H₂O after adjusting pH

Sterilize and cool to 60°C.

10 ## = Dissolve 1.660 g of Calcium Chloride Dihydrate in 950.000 ml of polished D-I H₂O. Then dissolve 4.629 of Ammonium Sulfate; 4.000 g of Potassium Phosphate Monobasic KH₂PO₄; 1.850 g of Magnesium Sulfate 7-H₂O, MgSO₄, 7H₂O; and 28.300 g of Potassium Nitrate into sequence. Bring up to volume with polished D-I H₂O.

= Dissolve 3.000 g of Boric Acid, 10.000 g of Manganous Sulfate Monohydrate, 0.250 g of Sodium Molybdate Dihydrate, and 0.750 g of Potassium Iodide in polished D-I H₂O in sequence. Bring up to volume with polished D-I H₂O.

- 5 #### = Dissolve 3.700 g of Disodium EDTA Dihydrate and 2.790 g of Ferrous Sulfate 7-Hydrate into D-I H₂O. Bring up to volume with D-I H₂O.

Total Volume (L) = 1.00

604 S

Ingredient	Amount	Unit
D-I H ₂ O	800.000	ml
CHU (N6) Basal Salts (SIGMA C-1416)	1.600	g
N6 Macronutrients 10X Stock ##	60.000	ml
Potassium Nitrate	1.680	g
B5H Minor Salts 1000X ###	0.600	ml
B5H Fe Na EDTA 100X ####	6.000	ml
Eriksson's Vitamin Mix (1000X SIGMA-1511)	0.400	ml
S & H Vitamin Mixture 100X Stock (S3766)	6.000	ml
Thiamine.HCL 0.4mg/ml	0.500	ml
L-Proline	1.980	g
Casein Hydrolysate (acid)	0.300	g
Sucrose	120.000	g
Glucose	0.600	g
2, 4-D 0.5mg/ml	1.600	ml
Gelrite @	2.000	g
Dicamba 1mg/ml #	1.200	ml
Silver Nitrate 2mg/ml #	1.700	ml

Directions:

@ = Add after bringing up to volume

5 # = Add after sterilizing and cooling to temp.

Dissolve ingredients in polished D-I H₂O in sequence

Adjust to pH 5.8

Bring up to volume with polished D-I H₂O after adjusting pH

Sterilize and cool to 60°C.

10 #### = Dissolve 1.660 g of Calcium Chloride Dihydrate in 950.000 ml of polished D-I H₂O. Then dissolve 4.629 of Ammonium Sulfate; 4.000 g of Potassium Phosphate Monobasic KH₂PO₄; 1.850 g of Magnesium Sulfate 7-H₂O, MgSO₄, 7H₂O; and 28.300 g of Potassium Nitrate into sequence. Bring up to volume with polished D-I H₂O.

= Dissolve 3.000 g of Boric Acid; 10.000 g of Manganous Sulfate Monohydrate; 0.250 g of Sodium Molybdate Dihydrate; and 0.750 g of Potassium Iodide in 950.000 ml of polished D-I H₂O in sequence. Bring up to volume with polished D-I H₂O.

- 5 ##### = Dissolve 3.700 g of Disodium EDTA Dihydrate and 2.790 g of Ferrous Sulfate 7-Hydrate into 950.000 ml of D-I H₂O. Bring up to volume with D-I H₂O.
Total Volume (L) = 1.00

272 V

Ingredient	Amount	Unit
D-I H ₂ O	950.000	ml
MS Salts (GIBCO 11117-074)	4.300	g
Myo-Inositol	0.100	g
MS Vitamins Stock Solution ##	5.000	ml
Sucrose	40.000	g
Bacto-Agar (@)	6.000	g

Directions:

(@) = Add after bringing up to volume

- 5 Dissolve ingredients in polished D-I H₂O in sequence

Adjust to pH 5.6

Bring up to volume with polished D-I H₂O after adjusting pH

Sterilize and cool to 60°C.

= Dissolve 0.100 g of Nicotinic Acid; 0.020 g of Thiamine.HCL; 0.100 g of

- 10 Pyridoxine.HCL; and 0.400 g of Glycine in 875.00 ml of polished D-I H₂O in sequence. Bring up to volume with polished D-I H₂O. Make in 400 ml portions. Thiamine.HCL & Pyridoxine.HCL are in Dark Desiccator. Store for one month, unless contamination or precipitation occur, then make fresh stock.

Total Volume (L) = 1.00

288 J

Ingredient	Amount	Unit
D-I H ₂ O	950.000	ml
MS Salts	4.300	g
Myo-Inositol	0.100	g
MS Vitamins Stock Solution ##	5.000	ml
Zeatin .5mg/ml	1.000	ml
Sucrose	60.000	g
Gelrite @	3.000	g
Indole Acetic Acid 0.5 mg/ml #	2.000	ml
.1mM Absissic Acid	1.000	ml
Bialaphos 1mg/ml #	3.000	ml

Directions:

@ = Add after bringing up to volume

- 5 Dissolve ingredients in polished D-I H₂O in sequence

Adjust to pH 5.6

Bring up to volume with polished D-I H₂O after adjusting pH

Sterilize and cool to 60°C.

Add 3.5g/L of Gelrite for cell biology.

- 10 ## = Dissolve 0.100 g of Nicotinic Acid; 0.020 g of Thiamine.HCL; 0.100 g of Pyridoxine.HCL; and 0.400 g of Glycine in 875.00 ml of polished D-I H₂O in sequence. Bring up to volume with polished D-I H₂O. Make in 400 ml portions. Thiamine.HCL & Pyridoxine.HCL are in Dark Desiccator. Store for one month, unless contamination or precipitation occur, then make fresh stock.

- 15 Total Volume (L) = 1.00

560 L

Ingredient	Amount	Unit
D-I Water, Filtered	950.000	ml
CHU (N6) Basal Salts (SIGMA C-1416)	4.000	g
Eriksson's Vitamin Mix (1000X SIGMA-1511)	0.400	ml
Thiamine.HCL 0.4mg/ml	1.250	ml
Sucrose	20.000	g
2, 4-D 0.5mg/ml	2.000	ml
L-Proline	2.880	g
Gelrite @	2.000	g
Silver Nitrate 2mg/ml #	4.250	ml

Directions:

@ = Add after bringing up to volume

5 # - Add after sterilizing and cooling to temp.

Dissolve ingredients in D-I H₂O in sequence

Adjust to pH 5.8 with KOH

Bring up to volume with D-I H₂O

Sterilize and cool to room temp.

10 Total Volume (L) = 1.00

560 R

Ingredient	Amount	Unit
D-I Water, Filtered	950.000	ml
CHU (N6) Basal Salts (SIGMA C-1416)	4.000	g
Eriksson's Vitamin Mix (1000X SIGMA-1511)	1.000	ml
Thiamine.HCL 0.4mg/ml	1.250	ml
Sucrose	30.000	g
2, 4-D 0.5mg/ml	4.000	ml
Gelrite @	3.000	g
Silver Nitrate 2mg/ml #	0.425	ml
Bialaphos 1mg/ml #	3.000	ml

Directions:

@ = Add after bringing up to volume

5 # = Add after sterilizing and cooling to temp.

Dissolve ingredients in D-I H₂O in sequence

Adjust to pH 5.8 with KOH

Bring up to volume with D-I H₂O

Sterilize and cool to room temp.

10 Total Volume (L) = 1.00

560 Y

Ingredient	Amount	Unit
D-I Water, Filtered	950.000	ml
CHU (N6) Basal Salts (SIGMA C-1416)	4.000	g
Eriksson's Vitamin Mix (1000X SIGMA-1511)	1.000	ml
Thiamine.HCL 0.4mg/ml	1.250	ml
Sucrose	120.000	g
2,4-D 0.5mg/ml	2.000	ml
L-Proline	2.880	g
Gelrite @	2.000	g
Silver Nitrate 2mg/ml #	4.250	ml

Directions:

@ = Add after bringing up to volume

5 # = Add after sterilizing and cooling to temp.

Dissolve ingredients in D-I H₂O in sequence

Adjust to pH 5.8 with KOH

Bring up to volume with D-I H₂O

Sterilize and cool to room temp.

10 ** Autoclave less time because of increased sucrose**

Total Volume (L) = 1.00

All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains.

All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically
5 and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

THAT WHICH IS CLAIMED:

1. A method for manipulating a metabolic pathway in a plant cell, said method comprising transforming a plant cell with at least one nucleotide sequence operably linked to a promoter that drives expression in a plant cell, wherein said nucleotide sequence is selected from the group consisting of:
- a) a nucleotide sequence encoding a limonene synthase protein,
 - b) a nucleotide sequence encoding a GPP synthase protein,
 - c) a nucleotide sequence encoding a carveol synthase protein,
 - 10 d) a nucleotide sequence encoding a S-linalool synthase protein,
 - e) a nucleotide sequence encoding a limonene synthase protein having the amino acid sequence set forth in SEQ ID NO:2,
 - f) a nucleotide sequence encoding a GPP synthase protein having the amino acid sequence set forth in SEQ ID NO:4,
 - 15 g) a nucleotide sequence encoding a carveol synthase protein having the amino acid sequence set forth in SEQ ID NO:6,
 - h) a nucleotide sequence encoding a S-linalool synthase protein having the amino acid sequence set forth in SEQ ID NO:8,
 - i) a nucleotide sequence encoding a limonene synthase protein as set forth in SEQ ID NO:1,
 - 20 j) a nucleotide sequence encoding a GPP synthase protein as set forth in SEQ ID NO:3,
 - k) a nucleotide sequence encoding a carveol synthase protein as set forth in SEQ ID NO:5,
 - 25 l) a nucleotide sequence encoding a S-linalool synthase protein as set forth in SEQ ID NO:7,
 - m) a nucleotide sequence corresponding to a portion of an antisense sequence for said nucleotide sequences of a, b, c, d, e, f, g, h, i, j, k or l.
- 30 2. The method of claim 1 wherein said plant cell is monocotyledonous.
3. The method of claim 2 wherein said monocot cell is a corn, wheat, rice, oat, rye or sorghum cell.

4. The method of claim 1 wherein said promoter is a constitutive promoter.

5. The method of claim 4 wherein said constitutive promoter is a ubiquitin promoter.

6. The method of claim 1 wherein said promoter is a tissue-specific promoter.

10

7. The method of claim 6 wherein said tissue-specific promoter is a root-specific promoter.

8. A method for manipulating a metabolic pathway in a plant cell, said method comprising transforming a plant cell with a first nucleotide sequence encoding a GPP synthase protein, and at least one nucleotide sequence which is selected from the group consisting of:

- a) a nucleotide sequence encoding a limonene synthase protein,
- b) a nucleotide sequence encoding a carveol synthase protein,
- 20 c) a nucleotide sequence encoding a S-linalool synthase protein,
- d) a nucleotide sequence encoding a limonene synthase protein having the amino acid sequence set forth in SEQ ID NO:2,
- e) a nucleotide sequence encoding a carveol synthase protein having the amino acid sequence set forth in SEQ ID NO:6,
- 25 f) a nucleotide sequence encoding a S-linalool synthase protein having the amino acid sequence set forth in SEQ ID NO:8,
- g) a nucleotide sequence encoding a limonene synthase protein as set forth in SEQ ID NO:1,
- h) A nucleotide sequence encoding a carveol synthase protein as forth in SEQ ID NO:5,
- 30 i) A nucleotide sequence encoding a S-linalool synthase protein as set forth in SEQ ID NO:7,

j) a nucleotide sequence corresponding to a portion of an antisense sequence for said nucleotide sequences of a, b, c, d, e, f, g, h or i; and
wherein said first and said second nucleotide sequence are operably linked to a promoter that drives expression in a plant cell.

5

9. The method of claim 8, wherein said plant cell is co-transformed.

10. A method for creating or enhancing resistance to insects in a plant, said method comprising manipulating a metabolic pathway in a plant cell according to the method of claim 8, wherein
said selected nucleotide sequence consists of a second nucleotide sequence encoding a limonene synthase protein, and
further comprising regenerating stably transformed plants producing effective amounts of limonene.

15

11. A method for creating or enhancing resistance to insects in a plant, said method comprising manipulating a metabolic pathway in a plant cell according to the method of claim 8, wherein

said selected nucleotide sequences consist of a second nucleotide sequence encoding a limonene synthase protein and
a third nucleotide sequence encoding a carveol synthase protein; and
further comprising regenerating stably transformed plants producing effective amounts of carveol.

25

12. A method for creating or enhancing resistance to insects in a plant, said method comprising manipulating a metabolic pathway in a plant cell according to the method of claim 8, wherein

said selected nucleotide sequence consists of a second nucleotide sequence encoding a S-linalool synthase protein, and
further comprising regenerating stably transformed plants producing effective amounts of S-linalool.

30

13. The method of any of the claims 10, 11, or 12 wherein said plant is monocotyledonous.

14. The method of claim 13 wherein said monocot plant is a corn plant
5 (Z. mays L.).

15. The method of claim 14 wherein said insect is larvae of *Diabrotica* spp.

10 16. A method of producing limonene synthase and GPP synthase in a plant cell, said method comprising transforming a first plant cell with a nucleotide sequence encoding a limonene synthase protein operably linked to a promoter that drives expression in a plant cell,

transforming a second plant cell with a nucleotide sequence encoding a
15 GPP synthase protein operably linked to a promoter that drives expression in a plant cell,

regenerating a first fertile transgenic plant from said first plant cell,
regenerating a second fertile transgenic plant from said second plant cell,
crossing said first and second fertile transgenic plants; and
20 recovering progeny from said cross which express limonene synthase and GPP synthase.

17. A method of producing limonene synthase, GPP synthase, and carveol synthase in a plant cell, said method comprising
25 transforming a first plant cell with a nucleotide sequence encoding a limonene synthase protein operably linked to a promoter that drives expression in a plant cell,

transforming a second plant cell with a nucleotide sequence encoding a
GPP synthase protein operably linked to a promoter that drives expression in a
30 plant cell,

transforming a third plant cell with a nucleotide sequence encoding a
carveol synthase protein operably linked to a promoter that drives expression in a
plant cell,

producing a first fertile transgenic plant from said first plant cell,
producing a second fertile transgenic plant from said second plant cell,
producing a third fertile transgenic plant from said third plant cell,
performing a first cross between said first and second fertile transgenic
5 plants,
recovering progeny from said first cross,
performing a second cross between said third fertile transgenic plant and
said progeny of said first cross; and
recovering progeny from said second cross which express limonene
10 synthase, GPP synthase and carveol synthase.

18. The method of claim 17, wherein said second plant cell is
transformed with a nucleotide sequence encoding a carveol synthase protein, and
said third plant cell is transformed with a nucleotide sequence encoding GPP
15 synthase.

19. A method of producing GPP synthase and S-linalool synthase in a
plant cell, said method comprising
transforming a first plant cell with a nucleotide sequence encoding a GPP
20 synthase protein operably linked to a promoter that drives expression in a plant
cell,
transforming a second plant cell with a nucleotide sequence encoding a S-
linalool synthase protein operably linked to a promoter that drives expression in a
plant cell,
25 producing a first fertile transgenic plant from said first plant cell,
producing a second fertile transgenic plant from said second plant cell,
crossing said first and second fertile transgenic plants; and
recovering progeny from said cross which express GPP synthase and S-
linalool synthase.

30

20. A transformed plant cell having a manipulated metabolic pathway
by having stably incorporated into its genome at least one nucleotide sequence
operably linked to a promoter that drives expression in a plant cell,

wherein said nucleotide sequence is selected from the group consisting of:

- a) a nucleotide sequence encoding a limonene synthase protein,
- b) a nucleotide sequence encoding a GPP synthase protein,
- c) a nucleotide sequence encoding a carveol synthase protein,
- 5 d) a nucleotide sequence encoding a S-linalool synthase protein,
- e) a nucleotide sequence encoding a limonene synthase protein having the amino acid sequence set forth in SEQ ID NO:2,
- f) a nucleotide sequence encoding a GPP synthase protein having the amino acid sequence set forth in SEQ ID NO:4,
- 10 g) a nucleotide sequence encoding a carveol synthase protein having the amino acid sequence set forth in SEQ ID NO:6,
- h) a nucleotide sequence encoding a S-linalool synthase protein having the amino acid sequence set forth in SEQ ID NO:8,
- i) a nucleotide sequence encoding a limonene synthase protein as set
- 15 forth in SEQ ID NO:1,
- j) a nucleotide sequence encoding a GPP synthase protein as set forth in SEQ ID NO:3,
- k) a nucleotide sequence encoding a carveol synthase protein as forth in SEQ ID NO:5,
- 20 l) a nucleotide sequence encoding a S-linalool synthase protein as set forth in SEQ ID NO:7,
- m) a nucleotide sequence corresponding to a portion of an antisense sequence for said nucleotide sequences of a, b, c, d, e, f, g, h, i, k or l.

- 25 21. A transformed plant having a manipulated metabolic pathway by having stably incorporated into its genome at least one nucleotide sequence operably linked to a promoter that drives expression in a plant cell,

wherein said nucleotide sequence is selected from the group consisting of:

- a) a nucleotide sequence encoding a limonene synthase protein,
- 30 b) a nucleotide sequence encoding a GPP synthase protein,
- c) a nucleotide sequence encoding a carveol synthase protein,
- d) a nucleotide sequence encoding a S-linalool synthase protein,

- e) a nucleotide sequence encoding a limonene synthase protein having the amino acid sequence set forth in SEQ ID NO:2,
- f) a nucleotide sequence encoding a GPP synthase protein having the amino acid sequence set forth in SEQ ID NO:4,
- 5 g) a nucleotide sequence encoding a carveol synthase protein having the amino acid sequence set forth in SEQ ID NO:6,
- h) a nucleotide sequence encoding a S-linalool synthase protein having the amino acid sequence set forth in SEQ ID NO:8,
- i) a nucleotide sequence encoding a limonene synthase protein as set
10 forth in SEQ ID NO:1,
- j) a nucleotide sequence encoding a GPP synthase protein as set forth in SEQ ID NO:3,
- k) a nucleotide sequence encoding a carveol synthase protein as set forth in SEQ ID NO:5,
- 15 l) a nucleotide sequence encoding a S-linalool synthase protein as set forth in SEQ ID NO:7,
- m) a nucleotide sequence corresponding to a portion of an antisense sequence for said nucleotide sequences of a, b, c, d, e, f, g, h, i, j, k or l.

20 22. The plant of claim 21 wherein said plant is monocotyledonous.

23. The plant of claim 22 wherein said monocot plant is a corn, wheat, rice, oat, rye or sorghum plant.

25 24. The plant of claim 21 wherein said promoter is a constitutive promoter.

25. The plant of claim 24 wherein said constitutive promoter is a ubiquitin promoter.

30

26. The plant of claim 21 wherein said promoter is a tissue-specific promoter.



27. The plant of claim 26 wherein said tissue-specific promoter is a root-specific promoter.

28. The plant of any of claims 24-27 wherein said plant is a dicot.

5

29. Seed of the plant of any of claims 21-27.

30. Seed of the plant of claim 28.








Figure 2





SEQUENCE LISTING

<110> EuClaire Meyer, Terry
Yalpani, Nasser

<120> Limonene and Other Downstream
Metabolites of Geranyl Pyrophosphate for Insect Control in
Plants

<130> 5718-65

<140> 09/172,339

<141> 1998-10-14

<150> 08/449,061

<151> 1995-05-24

<150> 08/153,544

<151> 1993-11-16

<150> 08/042,199

<151> 1993-04-02

<160> 8

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 2170

<212> DNA

<213> Mentha spicata

<220>

<221> misc_feature

<222> (0)...(0)

<223> Limonene Synthase

<221> CDS

<222> (29)...(1825)

<400> 1

```

agagagagag aggaaggaaa gattaatc atg gct ctc aaa gtg tta agt gtt      52
                               Met Ala Leu Lys Val Leu Ser Val
                               1                               5

gca act caa atg gcg att cct agc aac cta acg aca tgt ctt caa ccc      100
Ala Thr Gln Met Ala Ile Pro Ser Asn Leu Thr Thr Cys Leu Gln Pro
   10                               15                               20

tca cac ttc aaa tct tct cca aaa ctg tta tct agc act aac agt agt      148
Ser His Phe Lys Ser Ser Pro Lys Leu Leu Ser Ser Thr Asn Ser Ser
   25                               30                               35                               40

agt cgg tct cgc ctc cgt gtg tat tgc tcc tcc tcg caa ctc act act      196
Ser Arg Ser Arg Leu Arg Val Tyr Cys Ser Ser Ser Gln Leu Thr Thr
           45                               50                               55

```

gaa aga cga tcc gga aac tac aac cct tct cgt tgg gat gtc aac ttc Glu Arg Arg Ser Gly Asn Tyr Asn Pro Ser Arg Trp Asp Val Asn Phe 60 65 70	244
atc caa tcg ctt ctc agt gac tat aag gag gac aaa cac gtg att agg Ile Gln Ser Leu Leu Ser Asp Tyr Lys Glu Asp Lys His Val Ile Arg 75 80 85	292
gct tct gag ctg gtc act ttg gtg aag atg gaa ctg gag aaa gaa acg Ala Ser Glu Leu Val Thr Leu Val Lys Met Glu Leu Glu Lys Glu Thr 90 95 100	340
gat caa att cga caa ctt gag ttg atc gat gac ttg cag agg atg ggg Asp Gln Ile Arg Gln Leu Glu Leu Ile Asp Asp Leu Gln Arg Met Gly 105 110 115 120	388
ctg tcc gat cat ttc caa aat gag ttc aaa gaa atc ttg tcc tct ata Leu Ser Asp His Phe Gln Asn Glu Phe Lys Glu Ile Leu Ser Ser Ile 125 130 135	436
tat ctc gac cat cac tat tac aag aac cct ttt cca aaa gaa gaa agg Tyr Leu Asp His His Tyr Tyr Lys Asn Pro Phe Pro Lys Glu Glu Arg 140 145 150	484
gat ctc tac tcc aca tct ctt gca ttt agg ctc ctc aga gaa cat ggt Asp Leu Tyr Ser Thr Ser Leu Ala Phe Arg Leu Leu Arg Glu His Gly 155 160 165	532
ttt caa gtc gca caa gag gta ttc gat agt ttc aag aac gag gag ggt Phe Gln Val Ala Gln Glu Val Phe Asp Ser Phe Lys Asn Glu Glu Gly 170 175 180	580
gag ttc aaa gaa agc ctt agc gac gac acc aga gga ttg ttg caa ctg Glu Phe Lys Glu Ser Leu Ser Asp Asp Thr Arg Gly Leu Leu Gln Leu 185 190 195 200	628
tat gaa gct tcc ttt ctg ttg acg gaa ggc gaa acc acg ctc gag tca Tyr Glu Ala Ser Phe Leu Leu Thr Glu Gly Glu Thr Thr Leu Glu Ser 205 210 215	676
gcg agg gaa ttc gcc acc aaa ttt ttg gag gaa aaa gtg aac gag ggt Ala Arg Glu Phe Ala Thr Lys Phe Leu Glu Glu Lys Val Asn Glu Gly 220 225 230	724
ggt gtt gat ggc gac ctt tta aca aga atc gca tat tct ttg gac atc Gly Val Asp Gly Asp Leu Leu Thr Arg Ile Ala Tyr Ser Leu Asp Ile 235 240 245	772
cct ctt cat tgg agg att aaa agg cca aat gca cct gtg tgg atc gaa Pro Leu His Trp Arg Ile Lys Arg Pro Asn Ala Pro Val Trp Ile Glu 250 255 260	820
tgg tat agg aag agg ccc gac atg aat cca gta gtg ttg gag ctt gcc Trp Tyr Arg Lys Arg Pro Asp Met Asn Pro Val Val Leu Glu Leu Ala 265 270 275 280	868

ata ctc gac tta aat att gtt caa gca caa ttt caa gaa gag ctc aaa Ile Leu Asp Leu Asn Ile Val Gln Ala Gln Phe Gln Glu Glu Leu Lys 285 290 295	916
gaa tcc ttc agg tgg tgg aga aat act ggg ttt gtt gag aag ctg ccc Glu Ser Phe Arg Trp Trp Arg Asn Thr Gly Phe Val Glu Lys Leu Pro 300 305 310	964
ttc gca agg gat aga ctg gtg gaa tgc tac ttt tgg aat act ggg atc Phe Ala Arg Asp Arg Leu Val Glu Cys Tyr Phe Trp Asn Thr Gly Ile 315 320 325	1012
atc gag cca cgt cag cat gca agt gca agg ata atg atg ggc aaa gtc Ile Glu Pro Arg Gln His Ala Ser Ala Arg Ile Met Met Gly Lys Val 330 335 340	1060
aac gct ctg att acg gtg atc gat gat att tat gat gtc tat ggc acc Asn Ala Leu Ile Thr Val Ile Asp Asp Ile Tyr Asp Val Tyr Gly Thr 345 350 355 360	1108
tta gaa gaa ctc gaa caa ttc act gac ctc att cga aga tgg gat ata Leu Glu Glu Leu Glu Gln Phe Thr Asp Leu Ile Arg Arg Trp Asp Ile 365 370 375	1156
aac tca atc gac caa ctt ccc gat tac atg caa ctg tgc ttt ctt gca Asn Ser Ile Asp Gln Leu Pro Asp Tyr Met Gln Leu Cys Phe Leu Ala 380 385 390	1204
ctc aac aac ttc gtc gat gat aca tcg tac gat gtt atg aag gag aaa Leu Asn Asn Phe Val Asp Asp Thr Ser Tyr Asp Val Met Lys Glu Lys 395 400 405	1252
ggc gtc aac gtt ata ccc tac ctg cgg caa tcg tgg gtt gat ttg gcg Gly Val Asn Val Ile Pro Tyr Leu Arg Gln Ser Trp Val Asp Leu Ala 410 415 420	1300
gat aag tat atg gta gag gca cgg tgg ttc tac ggc ggg cac aaa cca Asp Lys Tyr Met Val Glu Ala Arg Trp Phe Tyr Gly Gly His Lys Pro 425 430 435 440	1348
agt ttg gaa gag tat ttg gag aac tca tgg cag tcg ata agt ggg ccc Ser Leu Glu Glu Tyr Leu Glu Asn Ser Trp Gln Ser Ile Ser Gly Pro 445 450 455	1396
tgt atg tta acg cac ata ttc ttc cga gta aca gat tcg ttc aca aag Cys Met Leu Thr His Ile Phe Phe Arg Val Thr Asp Ser Phe Thr Lys 460 465 470	1444
gag acc gtc gac agt ttg tac aaa tac cac gat tta gtt cgt tgg tca Glu Thr Val Asp Ser Leu Tyr Lys Tyr His Asp Leu Val Arg Trp Ser 475 480 485	1492
tcc ttc gtt ctg cgg ctt gct gat gat ttg gga acc tcg gtg gaa gag Ser Phe Val Leu Arg Leu Ala Asp Asp Leu Gly Thr Ser Val Glu Glu 490 495 500	1540
gtg agc aga ggg gat gtg ccg aaa tca ctt cag tgc tac atg agt gac	1588

Val	Ser	Arg	Gly	Asp	Val	Pro	Lys	Ser	Leu	Gln	Cys	Tyr	Met	Ser	Asp		
505					510					515					520		
tac	aat	gca	tcg	gag	gcg	gag	gcg	cgg	aag	cac	gtg	aaa	tgg	ctg	ata		1636
Tyr	Asn	Ala	Ser	Glu	Ala	Glu	Ala	Arg	Lys	His	Val	Lys	Trp	Leu	Ile		
				525					530					535			
gcg	gag	gtg	tgg	aag	aag	atg	aat	gcg	gag	agg	gtg	tcg	aag	gat	tct		1684
Ala	Glu	Val	Trp	Lys	Lys	Met	Asn	Ala	Glu	Arg	Val	Ser	Lys	Asp	Ser		
				540				545						550			
cca	ttc	ggc	aaa	gat	ttt	ata	gga	tgt	gca	gtt	gat	tta	gga	agg	atg		1732
Pro	Phe	Gly	Lys	Asp	Phe	Ile	Gly	Cys	Ala	Val	Asp	Leu	Gly	Arg	Met		
		555					560						565				
gcg	cag	ttg	atg	tac	cat	aat	gga	gat	ggg	cac	ggc	aca	caa	cac	cct		1780
Ala	Gln	Leu	Met	Tyr	His	Asn	Gly	Asp	Gly	His	Gly	Thr	Gln	His	Pro		
	570					575					580						
att	ata	cat	caa	caa	atg	acc	aga	acc	tta	ttc	gag	ccc	ttt	gca			1825
Ile	Ile	His	Gln	Gln	Met	Thr	Arg	Thr	Leu	Phe	Glu	Pro	Phe	Ala			
585					590					595							
tgagagatga	tgacgagcca	tcgtttactt	acttaaattc	taccaaagtt	tttcgaaggc												1885
atagttcgtg	atatttcaag	caccaataaa	taaggagaat	cggctcaaac	aaacgtggca												1945
tttgccacca	cgtgagcaca	aggagagatc	tgctgcgtgt	tatggatgaa	ctattcaatt												2005
tttatgcatg	taataattaa	gttcaagttc	aagagccttc	tgcataattta	actatgtatt												2065
tgaatttatc	gagtgtgatt	ttctgtcttt	ggcaacatat	atattttgtca	tatgtggcat												2125
cttattatga	tatcatacag	tgtttatgga	tgatatgata	ctatc													2170

<210> 2

<211> 599

<212> PRT

<213> Mentha spicata

<400> 2

Met	Ala	Leu	Lys	Val	Leu	Ser	Val	Ala	Thr	Gln	Met	Ala	Ile	Pro	Ser		
1				5					10					15			
Asn	Leu	Thr	Thr	Cys	Leu	Gln	Pro	Ser	His	Phe	Lys	Ser	Ser	Pro	Lys		
			20					25						30			
Leu	Leu	Ser	Ser	Thr	Asn	Ser	Ser	Ser	Arg	Ser	Arg	Leu	Arg	Val	Tyr		
		35				40						45					
Cys	Ser	Ser	Ser	Gln	Leu	Thr	Thr	Glu	Arg	Arg	Ser	Gly	Asn	Tyr	Asn		
	50					55					60						
Pro	Ser	Arg	Trp	Asp	Val	Asn	Phe	Ile	Gln	Ser	Leu	Leu	Ser	Asp	Tyr		
65					70				75					80			
Lys	Glu	Asp	Lys	His	Val	Ile	Arg	Ala	Ser	Glu	Leu	Val	Thr	Leu	Val		
				85				90						95			
Lys	Met	Glu	Leu	Glu	Lys	Glu	Thr	Asp	Gln	Ile	Arg	Gln	Leu	Glu	Leu		
		100					105						110				
Ile	Asp	Asp	Leu	Gln	Arg	Met	Gly	Leu	Ser	Asp	His	Phe	Gln	Asn	Glu		
	115					120					125						
Phe	Lys	Glu	Ile	Leu	Ser	Ser	Ile	Tyr	Leu	Asp	His	His	Tyr	Tyr	Lys		
	130					135				140							
Asn	Pro	Phe	Pro	Lys	Glu	Arg	Asp	Leu	Tyr	Ser	Thr	Ser	Leu	Ala			
145				150				155					160				
Phe	Arg	Leu	Leu	Arg	Glu	His	Gly	Phe	Gln	Val	Ala	Gln	Glu	Val	Phe		

<210> 3

<211> 1131
 <212> DNA
 <213> *Mentha spicata*

<220>
 <221> misc_feature
 <222> (0)...(0)
 <223> GFP Synthase

<221> CDS
 <222> (6)...(944)

<400> 3

tcaaa atg gcc att aat ctc tcc cat atc aac tcc aaa aca tgt ttc cct	50
Met Ala Ile Asn Leu Ser His Ile Asn Ser Lys Thr Cys Phe Pro	
1 5 10 15	
ctc aaa aca aga tct gat ctc agc cgt tct tct tcc gcg cgt tgc atg	98
Leu Lys Thr Arg Ser Asp Leu Ser Arg Ser Ser Ser Ala Arg Cys Met	
20 25 30	
cca act gcc gcc gct gcc gcc ttc ccc act atc gcc acc gcc gcc caa	146
Pro Thr Ala Ala Ala Ala Phe Pro Thr Ile Ala Thr Ala Ala Gln	
35 40 45	
agt cag ccg tac tgg gcc gcc atc gag gcc gac ata gag aga tac ctg	194
Ser Gln Pro Tyr Trp Ala Ala Ile Glu Ala Asp Ile Glu Arg Tyr Leu	
50 55 60	
aag aaa tcc atc aca ata agg ccg ccg gag aca gtt ttc ggg ccc atg	242
Lys Lys Ser Ile Thr Ile Arg Pro Pro Glu Thr Val Phe Gly Pro Met	
65 70 75	
cac cac ctc acc ttc gcc gcc cca gcc acc gcc gcc tcc acc cta tgc	290
His His Leu Thr Phe Ala Ala Pro Ala Thr Ala Ala Ser Thr Leu Cys	
80 85 90 95	
ttg gcg gcg tgc gag ctc gtc ggc ggc gac cga agc caa gcc atg gca	338
Leu Ala Ala Cys Glu Leu Val Gly Gly Asp Arg Ser Gln Ala Met Ala	
100 105 110	
gcc gcg gcg gcg atc cat ctc gtg cac gcg gca gcc tac gtc cac gag	386
Ala Ala Ala Ala Ile His Leu Val His Ala Ala Ala Tyr Val His Glu	
115 120 125	
cac ctc cct cta acc gac ggg tcg agg ccc gta tcc aag ccc gca atc	434
His Leu Pro Leu Thr Asp Gly Ser Arg Pro Val Ser Lys Pro Ala Ile	
130 135 140	
cag cac aag tac ggc ccg aac gtc gag ctc ctc acc gga gac ggg att	482
Gln His Lys Tyr Gly Pro Asn Val Glu Leu Leu Thr Gly Asp Gly Ile	
145 150 155	
gtc ccg ttc ggg ttt gag ttg ctg gcc ggg tca gtg gac ccg gcc cga	530
Val Pro Phe Gly Phe Glu Leu Leu Ala Gly Ser Val Asp Pro Ala Arg	
160 165 170 175	

aca gac gac ccg gat agg att ctg aga gtt ata ata gag atc agt cgg 578
 Thr Asp Asp Pro Asp Arg Ile Leu Arg Val Ile Ile Glu Ile Ser Arg
 180 185 190
 gcc ggc ggg ccg gag gga atg ata agc ggg ctg cat agg gaa gaa gaa 626
 Ala Gly Gly Pro Glu Gly Met Ile Ser Gly Leu His Arg Glu Glu Glu
 195 200 205
 att gtt gat gga aat acg agt tta gac ttc att gaa tat gtg tgc aag 674
 Ile Val Asp Gly Asn Thr Ser Leu Asp Phe Ile Glu Tyr Val Cys Lys
 210 215 220
 aaa aaa tac ggc gag atg cat gct tgc ggc gcg gct tgt gga gcc ata 722
 Lys Lys Tyr Gly Glu Met His Ala Cys Gly Ala Ala Cys Gly Ala Ile
 225 230 235
 ttg ggc ggc gca gcc gag gag gag att cag aag ctg agg aat ttc ggg 770
 Leu Gly Gly Ala Ala Glu Glu Glu Ile Gln Lys Leu Arg Asn Phe Gly
 240 245 250 255
 ctt tat caa gga act ctc aga gga atg atg gaa atg aaa aat tct cat 818
 Leu Tyr Gln Gly Thr Leu Arg Gly Met Met Glu Met Lys Asn Ser His
 260 265 270
 caa tta att gat gag aat ata att gga aaa ttg aaa gaa ttg gct ctc 866
 Gln Leu Ile Asp Glu Asn Ile Ile Gly Lys Leu Lys Glu Leu Ala Leu
 275 280 285
 gag gag ttg gga ggc ttc cac ggg aag aac gct gag ctg atg tgc agc 914
 Glu Glu Leu Gly Gly Phe His Gly Lys Asn Ala Glu Leu Met Ser Ser
 290 295 300
 ctt gta gcc gag ccg agc ctt tac gcg gct tagagctatt cggatccttc 964
 Leu Val Ala Glu Pro Ser Leu Tyr Ala Ala
 305 310
 attgcatttt catgcgacat cttcatattc atattgcata atatttttta agccagttat 1024
 ttttttatta tgaatttttt taactgttat tgatttcgaa aatactgaca atcatctaaa 1084
 ataaagtaaa tatagtaagg atgaaaaaaa aaaaaaaaaa aaaaaaa 1131

<210> 4

<211> 313

<212> PRT

<213> Mentha spicata

<400> 4

Met Ala Ile Asn Leu Ser His Ile Asn Ser Lys Thr Cys Phe Pro Leu
 1 5 10 15
 Lys Thr Arg Ser Asp Leu Ser Arg Ser Ser Ser Ala Arg Cys Met Pro
 20 25 30
 Thr Ala Ala Ala Ala Phe Pro Thr Ile Ala Thr Ala Ala Gln Ser
 35 40 45
 Gln Pro Tyr Trp Ala Ala Ile Glu Ala Asp Ile Glu Arg Tyr Leu Lys
 50 55 60
 Lys Ser Ile Thr Ile Arg Pro Pro Glu Thr Val Phe Gly Pro Met His
 65 70 75 80
 His Leu Thr Phe Ala Ala Pro Ala Thr Ala Ala Ser Thr Leu Cys Leu

			85					90				95			
Ala	Ala	Cys	Glu	Leu	Val	Gly	Gly	Asp	Arg	Ser	Gln	Ala	Met	Ala	Ala
			100					105					110		
Ala	Ala	Ala	Ile	His	Leu	Val	His	Ala	Ala	Ala	Tyr	Val	His	Glu	His
			115					120					125		
Leu	Pro	Leu	Thr	Asp	Gly	Ser	Arg	Pro	Val	Ser	Lys	Pro	Ala	Ile	Gln
			130				135					140			
His	Lys	Tyr	Gly	Pro	Asn	Val	Glu	Leu	Leu	Thr	Gly	Asp	Gly	Ile	Val
145					150					155				160	
Pro	Phe	Gly	Phe	Glu	Leu	Leu	Ala	Gly	Ser	Val	Asp	Pro	Ala	Arg	Thr
			165						170					175	
Asp	Asp	Pro	Asp	Arg	Ile	Leu	Arg	Val	Ile	Ile	Glu	Ile	Ser	Arg	Ala
			180					185					190		
Gly	Gly	Pro	Glu	Gly	Met	Ile	Ser	Gly	Leu	His	Arg	Glu	Glu	Glu	Ile
		195					200					205			
Val	Asp	Gly	Asn	Thr	Ser	Leu	Asp	Phe	Ile	Glu	Tyr	Val	Cys	Lys	Lys
	210				215						220				
Lys	Tyr	Gly	Glu	Met	His	Ala	Cys	Gly	Ala	Ala	Cys	Gly	Ala	Ile	Leu
225					230					235				240	
Gly	Gly	Ala	Ala	Glu	Glu	Glu	Ile	Gln	Lys	Leu	Arg	Asn	Phe	Gly	Leu
			245					250						255	
Tyr	Gln	Gly	Thr	Leu	Arg	Gly	Met	Met	Glu	Met	Lys	Asn	Ser	His	Gln
			260					265					270		
Leu	Ile	Asp	Glu	Asn	Ile	Ile	Gly	Lys	Leu	Lys	Glu	Leu	Ala	Leu	Glu
	275						280					285			
Glu	Leu	Gly	Gly	Phe	His	Gly	Lys	Asn	Ala	Glu	Leu	Met	Ser	Ser	Leu
	290					295					300				
Val	Ala	Glu	Pro	Ser	Leu	Tyr	Ala	Ala							
305					310										

<210> 5

<211> 1762

<212> DNA

<213> Mentha spicata

<220>

<221> misc_feature

<222> (0)...(0)

<223> Carveol Synthase

<221> CDS

<222> (20)...(1507)

<400> 5

aaaaaacwaa aaagaaacw atg gag ctc gac ctt ttg tgg gca att ata atc	52
Met Glu Leu Asp Leu Ser Ala Ile Ile Ile	
1 5 10	
ctt gtg gca acc tac atc gta tcc ctc cta atc aac caa tgg cga aaa	100
Leu Val Ala Thr Tyr Ile Val Ser Leu Leu Ile Asn Gln Trp Arg Lys	
15 20 25	
tgg aaa tcc caa caa aac cta cct ccg agc cct ccg aag ctg ccg gtg	148
Ser Lys Ser Gln Gln Asn Leu Pro Pro Ser Pro Pro Lys Leu Pro Val	
30 35 40	
atc ggc cac ctc cac ttc ctg tgg gga ggg ctt ccc cag cac gtg ttt	196

Ile	Gly	His	Leu	His	Phe	Leu	Trp	Gly	Gly	Leu	Pro	Gln	His	Val	Phe		
	45					50					55						
agg	agc	ata	gcc	cag	aag	tac	ggg	ccg	gtg	gcg	cac	gtg	cag	ctg	gga	244	
Arg	Ser	Ile	Ala	Gln	Lys	Tyr	Gly	Pro	Val	Ala	His	Val	Gln	Leu	Gly		
	60				65				70						75		
gaa	gtg	tac	tcg	gtg	gtg	ctg	tcg	tcg	gcg	gag	gca	gcg	ccg	cag	gcg	292	
Glu	Val	Tyr	Ser	Val	Val	Leu	Ser	Ser	Ala	Glu	Ala	Ala	Pro	Gln	Ala		
				80					85					90			
atg	aag	gtg	ctg	gac	ccg	aac	ttc	gcc	gac	cgg	ttc	gac	ggc	atc	ggg	340	
Met	Lys	Val	Leu	Asp	Pro	Asn	Phe	Ala	Asp	Arg	Phe	Asp	Gly	Ile	Gly		
			95					100					105				
tcc	agg	acc	atg	tgg	tac	gac	aaa	gat	gac	atc	atc	ttc	agc	cct	tac	388	
Ser	Arg	Thr	Met	Trp	Tyr	Asp	Lys	Asp	Asp	Ile	Ile	Phe	Ser	Pro	Tyr		
		110					115					120					
aac	gat	cac	tgg	cgc	cag	atg	cgg	agg	atc	tgc	gtg	aca	gag	ctg	ctg	436	
Asn	Asp	His	Trp	Arg	Gln	Met	Arg	Arg	Ile	Cys	Val	Thr	Glu	Leu	Leu		
	125					130					135						
agc	ccg	aag	aac	gtc	agg	tcc	ttc	ggg	tac	ata	agg	cag	gag	gag	atc	484	
Ser	Pro	Lys	Asn	Val	Arg	Ser	Phe	Gly	Tyr	Ile	Arg	Gln	Glu	Glu	Ile		
	140				145					150					155		
gag	cgc	ctc	atc	cgg	ctg	ctc	ggg	tcg	tcg	ggg	gga	gcg	ccg	gtc	gac	532	
Glu	Arg	Leu	Ile	Arg	Leu	Leu	Gly	Ser	Ser	Gly	Gly	Ala	Pro	Val	Asp		
				160					165					170			
gtg	acg	gag	gag	gtg	tcg	aag	atg	tcg	tgt	gtc	gtc	gtg	tgc	agg	gcg	580	
Val	Thr	Glu	Glu	Val	Ser	Lys	Met	Ser	Cys	Val	Val	Val	Cys	Arg	Ala		
				175				180					185				
gcg	ttc	ggg	agt	gtg	ctc	aag	gac	cag	ggt	tcg	ttg	gcg	gag	ttg	gtg	628	
Ala	Phe	Gly	Ser	Val	Leu	Lys	Asp	Gln	Gly	Ser	Leu	Ala	Glu	Leu	Val		
		190					195					200					
aag	gag	tcg	ctg	gca	ttg	gcg	tcc	ggg	ttt	gag	ctg	gcg	gat	ctc	tac	676	
Lys	Glu	Ser	Leu	Ala	Leu	Ala	Ser	Gly	Phe	Glu	Leu	Ala	Asp	Leu	Tyr		
	205					210					215						
cct	tcc	tca	tgg	ctc	ctc	aac	ctg	ctt	agc	ttg	aac	aag	tac	agg	ttg	724	
Pro	Ser	Ser	Trp	Leu	Leu	Asn	Leu	Leu	Ser	Leu	Asn	Lys	Tyr	Arg	Leu		
	220				225					230					235		
cag	agg	atg	cgc	cgc	cgc	ctc	gat	cac	atc	ctt	gat	ggg	ttc	ctg	gag	772	
Gln	Arg	Met	Arg	Arg	Arg	Leu	Asp	His	Ile	Leu	Asp	Gly	Phe	Leu	Glu		
				240					245					250			
gag	cat	agg	gag	aag	aag	agc	ggc	gac	ttt	gga	ggc	gag	gac	atc	gtc	820	
Glu	His	Arg	Glu	Lys	Lys	Ser	Gly	Asp	Phe	Gly	Gly	Glu	Asp	Ile	Val		
				255				260					265				
gac	gtt	ctt	ttc	agg	atg	cag	ccg	ggc	agc	gac	agc	aaa	att	ccc	att	868	
Asp	Val	Leu	Phe	Arg	Met	Gln	Pro	Gly	Ser	Asp	Ser	Lys	Ile	Pro	Ile		

270	275	280	
act tcc aat tgc atc aag ggt ttc att ttc gac acc ttc tcc gcg gga Thr Ser Asn Cys Ile Lys Gly Phe Ile Phe Asp Thr Phe Ser Ala Gly 285 290 295			916
gct gaa acg tct tgc acg acc atc tca tgg gcg ttg tgc gaa ctg atg Ala Glu Thr Ser Ser Thr Thr Ile Ser Trp Ala Leu Ser Glu Leu Met 300 305 310 315			964
agg aat ccg gcg aag atg gcc aag gtg cag gcg gag gta aga gag gcg Arg Asn Pro Ala Lys Met Ala Lys Val Gln Ala Glu Val Arg Glu Ala 320 325 330			1012
ctc aag gga aag aca gtc gtg gat ttg agc gag gtg caa gag cta aaa Leu Lys Gly Lys Thr Val Val Asp Leu Ser Glu Val Gln Glu Leu Lys 335 340 345			1060
tac ctg aga tgc gtg tta aag gag act ctg agg ctg cac cct ccc ttt Tyr Leu Arg Ser Val Leu Lys Glu Thr Leu Arg Leu His Pro Pro Phe 350 355 360			1108
cca tta atc cca aga caa tcc agg gaa gaa tgc gag gtt aac ggg tac Pro Leu Ile Pro Arg Gln Ser Arg Glu Glu Cys Glu Val Asn Gly Tyr 365 370 375			1156
acg att ccg gcc aaa act aga atc ttc atc aac gtc tgg gct atc gga Thr Ile Pro Ala Lys Thr Arg Ile Phe Ile Asn Val Trp Ala Ile Gly 380 385 390 395			1204
agg gat ccc caa tac tgg gaa gat ccc gac acc ttc cgc cct gag aga Arg Asp Pro Gln Tyr Trp Glu Asp Pro Asp Thr Phe Arg Pro Glu Arg 400 405 410			1252
ttc gat gag gtt tcc agg gat ttc atg gga aac gat ttc gag ttc atc Phe Asp Glu Val Ser Arg Asp Phe Met Gly Asn Asp Phe Glu Phe Ile 415 420 425			1300
cca ttc ggg gcg ggt cga aga atc tgc ccc ggt tta cat ttc ggg ctg Pro Phe Gly Ala Gly Arg Arg Ile Cys Pro Gly Leu His Phe Gly Leu 430 435 440			1348
gca aat gtt gag atc cca ttg gcg caa ctg ctc tac cac ttc gac tgg Ala Asn Val Glu Ile Pro Leu Ala Gln Leu Leu Tyr His Phe Asp Trp 445 450 455			1396
aaa ttg cca caa gga atg act gat gcc gac ttg gca ctg acg gag acc Lys Leu Pro Gln Gly Met Thr Asp Ala Asp Leu Ala Leu Thr Glu Thr 460 465 470 475			1444
cca ggt ctt tct ggg cca aaa aag aaa aat gtt tgc ttg gtt ccc aca Pro Gly Leu Ser Gly Pro Lys Lys Lys Asn Val Cys Leu Val Pro Thr 480 485 490			1492
ctc tat aaa agt cct taaccactaa gaagtttagca taataagaca tctaaaattg Leu Tyr Lys Ser Pro 495			1547

tcataatcat ctaattattg ttacacttct tctatcatgt catttttgaga agtgtcttat 1607
 agaggtggcc acggttccgg ttccagttcg gaagcggaac cgaaccatca gttacgggtc 1667
 tcagcaagaa gcgaaccgtc ccgccccccc tactgtgttt gagatataaa acacataaaa 1727
 taaaataaaa aaaacgctat ttttttttaa aaaaa 1762

<210> 6

<211> 496

<212> PRT

<213> *Mentha spicata*

<400> 6

Met	Glu	Leu	Asp	Leu	Leu	Ser	Ala	Ile	Ile	Ile	Leu	Val	Ala	Thr	Tyr	1	5	10	15
Ile	Val	Ser	Leu	Leu	Ile	Asn	Gln	Trp	Arg	Lys	Ser	Lys	Ser	Gln	Gln	20	25	30	
Asn	Leu	Pro	Pro	Ser	Pro	Pro	Lys	Leu	Pro	Val	Ile	Gly	His	Leu	His	35	40	45	
Phe	Leu	Trp	Gly	Gly	Leu	Pro	Gln	His	Val	Phe	Arg	Ser	Ile	Ala	Gln	50	55	60	
Lys	Tyr	Gly	Pro	Val	Ala	His	Val	Gln	Leu	Gly	Glu	Val	Tyr	Ser	Val	65	70	75	80
Val	Leu	Ser	Ser	Ala	Glu	Ala	Ala	Pro	Gln	Ala	Met	Lys	Val	Leu	Asp	85	90	95	
Pro	Asn	Phe	Ala	Asp	Arg	Phe	Asp	Gly	Ile	Gly	Ser	Arg	Thr	Met	Trp	100	105	110	
Tyr	Asp	Lys	Asp	Asp	Ile	Ile	Phe	Ser	Pro	Tyr	Asn	Asp	His	Trp	Arg	115	120	125	
Gln	Met	Arg	Arg	Ile	Cys	Val	Thr	Glu	Leu	Leu	Ser	Pro	Lys	Asn	Val	130	135	140	
Arg	Ser	Phe	Gly	Tyr	Ile	Arg	Gln	Glu	Glu	Ile	Glu	Arg	Leu	Ile	Arg	145	150	155	160
Leu	Leu	Gly	Ser	Ser	Gly	Gly	Ala	Pro	Val	Asp	Val	Thr	Glu	Glu	Val	165	170	175	
Ser	Lys	Met	Ser	Cys	Val	Val	Val	Cys	Arg	Ala	Ala	Phe	Gly	Ser	Val	180	185	190	
Leu	Lys	Asp	Gln	Gly	Ser	Leu	Ala	Glu	Leu	Val	Lys	Glu	Ser	Leu	Ala	195	200	205	
Leu	Ala	Ser	Gly	Phe	Glu	Leu	Ala	Asp	Leu	Tyr	Pro	Ser	Ser	Trp	Leu	210	215	220	
Leu	Asn	Leu	Leu	Ser	Leu	Asn	Lys	Tyr	Arg	Leu	Gln	Arg	Met	Arg	Arg	225	230	235	240
Arg	Leu	Asp	His	Ile	Leu	Asp	Gly	Phe	Leu	Glu	Glu	His	Arg	Glu	Lys	245	250	255	
Lys	Ser	Gly	Asp	Phe	Gly	Gly	Glu	Asp	Ile	Val	Asp	Val	Leu	Phe	Arg	260	265	270	
Met	Gln	Pro	Gly	Ser	Asp	Ser	Lys	Ile	Pro	Ile	Thr	Ser	Asn	Cys	Ile	275	280	285	
Lys	Gly	Phe	Ile	Phe	Asp	Thr	Phe	Ser	Ala	Gly	Ala	Glu	Thr	Ser	Ser	290	295	300	
Thr	Thr	Ile	Ser	Trp	Ala	Leu	Ser	Glu	Leu	Met	Arg	Asn	Pro	Ala	Lys	305	310	315	320
Met	Ala	Lys	Val	Gln	Ala	Glu	Val	Arg	Glu	Ala	Leu	Lys	Gly	Lys	Thr	325	330	335	
Val	Val	Asp	Leu	Ser	Glu	Val	Gln	Glu	Leu	Lys	Tyr	Leu	Arg	Ser	Val	340	345	350	
Leu	Lys	Glu	Thr	Leu	Arg	Leu	His	Pro	Pro	Phe	Pro	Leu	Ile	Pro	Arg				

	355		360		365										
Gln	Ser	Arg	Glu	Glu	Cys	Glu	Val	Asn	Gly	Tyr	Thr	Ile	Pro	Ala	Lys
	370					375					380				
Thr	Arg	Ile	Phe	Ile	Asn	Val	Trp	Ala	Ile	Gly	Arg	Asp	Pro	Gln	Tyr
385					390					395					400
Trp	Glu	Asp	Pro	Asp	Thr	Phe	Arg	Pro	Glu	Arg	Phe	Asp	Glu	Val	Ser
				405					410					415	
Arg	Asp	Phe	Met	Gly	Asn	Asp	Phe	Glu	Phe	Ile	Pro	Phe	Gly	Ala	Gly
			420					425					430		
Arg	Arg	Ile	Cys	Pro	Gly	Leu	His	Phe	Gly	Leu	Ala	Asn	Val	Glu	Ile
	435						440						445		
Pro	Leu	Ala	Gln	Leu	Leu	Tyr	His	Phe	Asp	Trp	Lys	Leu	Pro	Gln	Gly
	450					455					460				
Met	Thr	Asp	Ala	Asp	Leu	Ala	Leu	Thr	Glu	Thr	Pro	Gly	Leu	Ser	Gly
465					470					475					480
Pro	Lys	Lys	Lys	Asn	Val	Cys	Leu	Val	Pro	Thr	Leu	Tyr	Lys	Ser	Pro
				485					490					495	

<211> 7
 <212> 2681
 <213> DNA
 <214> Clarkia breweri

<220>
 <221> misc_feature
 <222> (0)...(0)
 <223> S-Linalool Synthase

<221> CDS
 <222> (28)...(2637)

<400> 7
 aaccaaacca ccttaaaca gacaacc atg cag ctc ata aca aat ttc tcc tca 54
 Met Gln Leu Ile Thr Asn Phe Ser Ser
 1 5

tca tca tca gaa ttg cag ttt ctt gtg gat aag gtt aag aga gaa tca 102
 Ser Ser Ser Glu Leu Gln Phe Leu Val Asp Lys Val Lys Arg Glu Ser
 10 15 20 25

ttg tct tct tca tca tct aat act cag aat ttg ttt ctc tca act tca 150
 Leu Ser Ser Ser Ser Ser Asn Thr Gln Asn Leu Phe Leu Ser Thr Ser
 30 35 40

cct tat gac act gct tgg ctc gcc ctt atc cct cat cct cat cat cac 198
 Pro Tyr Asp Thr Ala Trp Leu Ala Leu Ile Pro His Pro His His His
 45 50 55

cat cac cat ggc cga ccc atg ttt gaa aaa tgt ctg caa tgg att ctc 246
 His His His Gly Arg Pro Met Phe Glu Lys Cys Leu Gln Trp Ile Leu
 60 65 70

cat aac cag aca cca caa ggt ttc tgg gca gca gct ggt gac aat att 294
 His Asn Gln Thr Pro Gln Gly Phe Trp Ala Ala Ala Gly Asp Asn Ile
 75 80 85

tcc gac acc gac gat gac gtc acc ctg gat tgt ctt cta tca acc ttg 342

Ser 90	Asp	Thr	Asp	Asp	Asp	Val	Thr	Leu	Asp	Cys 100	Leu	Leu	Ser	Thr	Leu 105	
gct	tgc	tta	gtt	gca	ctc	aaa	agg	tgg	cag	ctt	gct	ccc	gac	atg	att	390
Ala	Cys	Leu	Val	Ala	Leu	Lys	Arg	Trp	Gln	Leu	Ala	Pro	Asp	Met	Ile	
				110					115					120		
cat	aaa	gga	ttg	gaa	ttt	gta	aat	aga	aac	aca	gag	aga	ctt	gta	atg	438
His	Lys	Gly	Leu	Glu	Phe	Val	Asn	Arg	Asn	Thr	Glu	Arg	Leu	Val	Met	
			125					130					135			
aag	cag	aag	ccg	agc	gac	gtt	cct	cgt	tgg	ttc	acc	atc	atg	ttc	ccg	486
Lys	Gln	Lys	Pro	Ser	Asp	Val	Pro	Arg	Trp	Phe	Thr	Ile	Met	Phe	Pro	
		140					145					150				
gcg	atg	ctc	gag	ctt	gcc	gga	gct	tcc	agt	ctc	cga	gtc	gat	ttc	agc	534
Ala	Met	Leu	Glu	Leu	Ala	Gly	Ala	Ser	Ser	Leu	Arg	Val	Asp	Phe	Ser	
	155					160					165					
gag	aat	ctt	aac	aga	atc	ttg	gtg	gaa	cta	tct	caa	aat	agg	gat	gat	582
Glu	Asn	Leu	Asn	Arg	Ile	Leu	Val	Glu	Leu	Ser	Gln	Asn	Arg	Asp	Asp	
170					175					180					185	
att	ctc	aca	agg	gag	gaa	gtt	gat	gag	aag	aag	caa	tac	tca	cca	ttg	630
Ile	Leu	Thr	Arg	Glu	Glu	Val	Asp	Glu	Lys	Lys	Gln	Tyr	Ser	Pro	Leu	
				190					195					200		
cta	cta	ttt	cta	gaa	gca	ttg	cct	gca	caa	tcc	tat	gac	aat	gat	gtt	678
Leu	Leu	Phe	Leu	Glu	Ala	Leu	Prc	Ala	Gln	Ser	Tyr	Asp	Asn	Asp	Val	
			205					210					215			
cta	aag	caa	att	ata	gac	aag	aac	ttg	agc	aat	gat	ggc	tct	tta	ttg	726
Leu	Lys	Gln	Ile	Ile	Asp	Lys	Asn	Leu	Ser	Asn	Asp	Gly	Ser	Leu	Leu	
		220					225					230				
caa	tcg	cct	tct	gct	aca	gca	aga	gca	tac	atg	ata	aca	gga	aat	acc	774
Gln	Ser	Prc	Ser	Ala	Thr	Ala	Arg	Ala	Tyr	Met	Ile	Thr	Gly	Asn	Thr	
	235					240					245					
aga	tgc	tta	tcg	tat	cta	cac	tct	tta	aca	aat	agc	tgc	tct	aat	gga	822
Arg	Cys	Leu	Ser	Tyr	Leu	His	Ser	Leu	Thr	Asn	Ser	Cys	Ser	Asn	Gly	
250					255					260					265	
gga	gta	cca	tca	ttc	tat	cct	gtt	gac	gac	gac	ctc	cat	gat	ctt	gtc	870
Gly	Val	Pro	Ser	Phe	Tyr	Pro	Val	Asp	Asp	Asp	Leu	His	Asp	Leu	Val	
				270				275						280		
atg	gtg	aat	caa	ctg	aca	agg	tcg	ggc	ttg	act	gaa	cat	ctc	atc	ccg	918
Met	Val	Asn	Gln	Leu	Thr	Arg	Ser	Gly	Leu	Thr	Glu	His	Leu	Ile	Pro	
			285					290					295			
gag	att	gac	cac	ctt	cta	ctc	aaa	gtt	caa	aag	aac	tac	aaa	tac	aaa	966
Glu	Ile	Asp	His	Leu	Leu	Leu	Lys	Val	Gln	Lys	Asn	Tyr	Lys	Tyr	Lys	
		300					305					310				
aaa	gca	tca	cca	aaa	tca	ttg	tat	agc	att	gct	gcg	gaa	cta	tac	agg	1014
Lys	Ala	Ser	Pro	Lys	Ser	Leu	Tyr	Ser	Ile	Ala	Ala	Glu	Leu	Tyr	Arg	

315	320	325	
gat tca tta gca ttt tgg ttg ctt cga gtc aat aat cac tgg gta tca Asp Ser Leu Ala Phe Trp Leu Leu Arg Val Asn Asn His Trp Val Ser 330 335 340 345			1062
cca tca att ttt tgt tgg ttt tta gat gac gac gaa atc cgt gat cac Pro Ser Ile Phe Cys Trp Phe Leu Asp Asp Asp Glu Ile Arg Asp His 350 355 360			1110
atc gaa aca aac tac gag gaa ttt gct gcc gtg ctt ctt aat gtg tat Ile Glu Thr Asn Tyr Glu Glu Phe Ala Ala Val Leu Leu Asn Val Tyr 365 370 375			1158
cga gct acc gat ctt atg ttc tcc ggc gaa gtc caa ctt gtc gaa gca Arg Ala Thr Asp Leu Met Phe Ser Gly Glu Val Gln Leu Val Glu Ala 380 385 390			1206
aga tct ttc gct acc aag aat ctt gag aaa ata tta gca aca gga aac Arg Ser Phe Ala Thr Lys Asn Leu Glu Lys Ile Leu Ala Thr Gly Asn 395 400 405			1254
ata cat aaa act aat gca gat atc tca tct agt ttg cat aag atg atc Ile His Lys Thr Asn Ala Asp Ile Ser Ser Ser Leu His Lys Met Ile 410 415 420 425			1302
gaa cac gaa cta aga gtt cct tgg acc gca aga atg gac cat gtt gaa Glu His Glu Leu Arg Val Pro Trp Thr Ala Arg Met Asp His Val Glu 430 435 440			1350
aat cga att tgg atc gaa gaa ata gct tcc agt gct tta tgg ttt gga Asn Arg Ile Trp Ile Glu Glu Ile Ala Ser Ser Ala Leu Trp Phe Gly 445 450 455			1398
aaa tca tcc tac ctt agg tta tct tgc ttt cac aag atg agt tta cag Lys Ser Ser Tyr Leu Arg Leu Ser Cys Phe His Lys Met Ser Leu Gln 460 465 470			1446
caa ctc gcg gtg aaa aat tat acg ctt cga caa ttg gtt tac cga gac Gln Leu Ala Val Lys Asn Tyr Thr Leu Arg Gln Leu Val Tyr Arg Asp 475 480 485			1494
gag ctt gcg gaa gtt gag agg tgg tct aaa gaa aga ggg cta tgt gac Glu Leu Ala Glu Val Glu Arg Trp Ser Lys Glu Arg Gly Leu Cys Asp 490 495 500 505			1542
atg gga ttt tgt aga gag aaa acc ggg tat tgt tac tac gca ttt gcg Met Gly Phe Cys Arg Glu Lys Thr Gly Tyr Cys Tyr Tyr Ala Phe Ala 510 515 520			1590
gca agt act tgt ctg ccg tgg agt tcc gac gtg agg ctg gtc ctg acc Ala Ser Thr Cys Leu Pro Trp Ser Ser Asp Val Arg Leu Val Leu Thr 525 530 535			1638
aag gcg gca gtt gtc att aca gtg gcc gat gat ttc ttt gat gtc gaa Lys Ala Ala Val Val Ile Thr Val Ala Asp Asp Phe Phe Asp Val Glu 540 545 550			1686

gga tct atg gtt gat ctc gaa aaa tta acg gat gca gtt cgg agg tgg Gly Ser Met Val Asp Leu Glu Lys Leu Thr Asp Ala Val Arg Arg Trp 555 560 565	1734
gat gcg gaa ggg tta ggc agc cac agc aag aca ata ttt gaa gcc ctg Asp Ala Glu Gly Leu Gly Ser His Ser Lys Thr Ile Phe Glu Ala Leu 570 575 580 585	1782
gat gat ctt gta aat gaa gtt aga ctc aag tgt ttc caa caa aat gga Asp Asp Leu Val Asn Glu Val Arg Leu Lys Cys Phe Gln Gln Asn Gly 590 595 600	1830
caa gac atc aaa aac aat ctc caa caa tta tgg tat gaa aca ttc cat Gln Asp Ile Lys Asn Asn Leu Gln Gln Leu Trp Tyr Glu Thr Phe His 605 610 615	1878
tca tgg ctt atg gaa gct aag tgg gga aag ggg tta aca agt aaa cca Ser Trp Leu Met Glu Ala Lys Trp Gly Lys Gly Leu Thr Ser Lys Pro 620 625 630	1926
tct gta gat gtg tat ctt gga aat gca atg aca tcc ata gca gct cac Ser Val Asp Val Tyr Leu Gly Asn Ala Met Thr Ser Ile Ala Ala His 635 640 645	1974
acc atg gtc ctt aca gca tcc tgt ctt cta ggt ccc ggt ttc ccg gtt Thr Met Val Leu Thr Ala Ser Cys Leu Leu Gly Pro Gly Phe Pro Val 650 655 660 665	2022
cac caa cta tgg tcg caa agg cgc cac cag gac att aca tcc ttg ctc His Gln Leu Trp Ser Gln Arg Arg His Gln Asp Ile Thr Ser Leu Leu 670 675 680	2070
atg gtc ttg act cgc ttg cta aat gac att caa tcc tac ttg aaa gaa Met Val Leu Thr Arg Leu Leu Asn Asp Ile Gln Ser Tyr Leu Lys Glu 685 690 695	2118
gaa gac gaa gga aaa ata aac tat gta tgg atg tac atg atc gag aac Glu Asp Glu Gly Lys Ile Asn Tyr Val Trp Met Tyr Met Ile Glu Asn 700 705 710	2166
aat caa gcg tcg ata gat gac tcg gtt cga cac gtc cag acg ata atc Asn Gln Ala Ser Ile Asp Asp Ser Val Arg His Val Gln Thr Ile Ile 715 720 725	2214
aat gta aaa aag caa gaa ttc atc caa cgt gtt cta tcg gat caa cat Asn Val Lys Lys Gln Glu Phe Ile Gln Arg Val Leu Ser Asp Gln His 730 735 740 745	2262
tgc aat ctc cca aag tca ttc aag cag ctc cat ttc tcc tgc ctc aaa Cys Asn Leu Pro Lys Ser Phe Lys Gln Leu His Phe Ser Cys Leu Lys 750 755 760	2310
gta ttc aac atg ttc ttc aac tcc tcc aac att ttc gac act gat acc Val Phe Asn Met Phe Phe Asn Ser Ser Asn Ile Phe Asp Thr Asp Thr 765 770 775	2358

gac ctt ctt ctt gac att cac gaa gct ttt gtt tct cca cca caa gtt 2406
 Asp Leu Leu Leu Asp Ile His Glu Ala Phe Val Ser Pro Pro Gln Val
 780 785 790

ccc aaa ttc aaa ccc cac atc aag cca cct cat cag ctt cca gca aca 2454
 Pro Lys Phe Lys Pro His Ile Lys Pro Pro His Gln Leu Pro Ala Thr
 795 800 805

ctt cag cca cct cat cag ccc caa caa ata atg gtc aat aag aag aag 2502
 Leu Gln Pro Pro His Gln Pro Gln Gln Ile Met Val Asn Lys Lys Lys
 810 815 820 825

gtg gaa atg gtt tac aaa agc tat cat cat cca ttc aag gtt ttc acc 2550
 Val Glu Met Val Tyr Lys Ser Tyr His His Pro Phe Lys Val Phe Thr
 830 835 840

ttg cag aag aaa caa agt tcg gga cat ggt aca atg aat cca agg gct 2598
 Leu Gln Lys Lys Gln Ser Ser Gly His Gly Thr Met Asn Pro Arg Ala
 845 850 855

agt atc tta gca gga ccc aac atc aaa cta tgt ttc agt taacgaatac 2647
 Ser Ile Leu Ala Gly Pro Asn Ile Lys Leu Cys Phe Ser
 860 865 870

actaccttgt tattagaaga tgtcaccagt ttcc 2681

<210> 8
 <211> 870
 <212> PRT
 <213> Clarkia breweri

<400> 8
 Met Gln Leu Ile Thr Asn Phe Ser Ser Ser Ser Ser Glu Leu Gln Phe
 1 5 10 15
 Leu Val Asp Lys Val Lys Arg Glu Ser Leu Ser Ser Ser Ser Ser Asn
 20 25 30
 Thr Gln Asn Leu Phe Leu Ser Thr Ser Pro Tyr Asp Thr Ala Trp Leu
 35 40 45
 Ala Leu Ile Pro His Pro His His His His His Gly Arg Pro Met
 50 55 60
 Phe Glu Lys Cys Leu Gln Trp Ile Leu His Asn Gln Thr Pro Gln Gly
 65 70 75 80
 Phe Trp Ala Ala Ala Gly Asp Asn Ile Ser Asp Thr Asp Asp Asp Val
 85 90 95
 Thr Leu Asp Cys Leu Leu Ser Thr Leu Ala Cys Leu Val Ala Leu Lys
 100 105 110
 Arg Trp Gln Leu Ala Pro Asp Met Ile His Lys Gly Leu Glu Phe Val
 115 120 125
 Asn Arg Asn Thr Glu Arg Leu Val Met Lys Gln Lys Pro Ser Asp Val
 130 135 140
 Pro Arg Trp Phe Thr Ile Met Phe Pro Ala Met Leu Glu Leu Ala Gly
 145 150 155 160
 Ala Ser Ser Leu Arg Val Asp Phe Ser Glu Asn Leu Asn Arg Ile Leu
 165 170 175
 Val Glu Leu Ser Gln Asn Arg Asp Asp Ile Leu Thr Arg Glu Glu Val
 180 185 190
 Asp Glu Lys Lys Gln Tyr Ser Pro Leu Leu Leu Phe Leu Glu Ala Leu

195	200	205
Pro Ala Gln Ser Tyr Asp Asn Asp Val Leu Lys Gln Ile Ile Asp Lys		
210	215	220
Asn Leu Ser Asn Asp Gly Ser Leu Leu Gln Ser Pro Ser Ala Thr Ala		
225	230	235
Arg Ala Tyr Met Ile Thr Gly Asn Thr Arg Cys Leu Ser Tyr Leu His		
245	250	255
Ser Leu Thr Asn Ser Cys Ser Asn Gly Gly Val Pro Ser Phe Tyr Pro		
260	265	270
Val Asp Asp Asp Leu His Asp Leu Val Met Val Asn Gln Leu Thr Arg		
275	280	285
Ser Gly Leu Thr Glu His Leu Ile Pro Glu Ile Asp His Leu Leu Leu		
290	295	300
Lys Val Gln Lys Asn Tyr Lys Tyr Lys Lys Ala Ser Pro Lys Ser Leu		
305	310	315
Tyr Ser Ile Ala Ala Glu Leu Tyr Arg Asp Ser Leu Ala Phe Trp Leu		
325	330	335
Leu Arg Val Asn Asn His Trp Val Ser Pro Ser Ile Phe Cys Trp Phe		
340	345	350
Leu Asp Asp Asp Glu Ile Arg Asp His Ile Glu Thr Asn Tyr Glu Glu		
355	360	365
Phe Ala Ala Val Leu Leu Asn Val Tyr Arg Ala Thr Asp Leu Met Phe		
370	375	380
Ser Gly Glu Val Gln Leu Val Glu Ala Arg Ser Phe Ala Thr Lys Asn		
385	390	395
Leu Glu Lys Ile Leu Ala Thr Gly Asn Ile His Lys Thr Asn Ala Asp		
405	410	415
Ile Ser Ser Ser Leu His Lys Met Ile Glu His Glu Leu Arg Val Pro		
420	425	430
Trp Thr Ala Arg Met Asp His Val Glu Asn Arg Ile Trp Ile Glu Glu		
435	440	445
Ile Ala Ser Ser Ala Leu Trp Phe Gly Lys Ser Ser Tyr Leu Arg Leu		
450	455	460
Ser Cys Phe His Lys Met Ser Leu Gln Gln Leu Ala Val Lys Asn Tyr		
465	470	475
Thr Leu Arg Gln Leu Val Tyr Arg Asp Glu Leu Ala Glu Val Glu Arg		
485	490	495
Trp Ser Lys Glu Arg Gly Leu Cys Asp Met Gly Phe Cys Arg Glu Lys		
500	505	510
Thr Gly Tyr Cys Tyr Tyr Ala Phe Ala Ala Ser Thr Cys Leu Pro Trp		
515	520	525
Ser Ser Asp Val Arg Leu Val Leu Thr Lys Ala Ala Val Val Ile Thr		
530	535	540
Val Ala Asp Asp Phe Phe Asp Val Glu Gly Ser Met Val Asp Leu Glu		
545	550	555
Lys Leu Thr Asp Ala Val Arg Arg Trp Asp Ala Glu Gly Leu Gly Ser		
565	570	575
His Ser Lys Thr Ile Phe Glu Ala Leu Asp Asp Leu Val Asn Glu Val		
580	585	590
Arg Leu Lys Cys Phe Gln Gln Asn Gly Gln Asp Ile Lys Asn Asn Leu		
595	600	605
Gln Gln Leu Trp Tyr Glu Thr Phe His Ser Trp Leu Met Glu Ala Lys		
610	615	620
Trp Gly Lys Gly Leu Thr Ser Lys Pro Ser Val Asp Val Tyr Leu Gly		
625	630	635
Asn Ala Met Thr Ser Ile Ala Ala His Thr Met Val Leu Thr Ala Ser		
645	650	655


Cys Leu Leu Gly Pro Gly Phe Pro Val His Gln Leu Trp Ser Gln Arg
 660 665 670
 Arg His Gln Asp Ile Thr Ser Leu Leu Met Val Leu Thr Arg Leu Leu
 675 680 685
 Asn Asp Ile Gln Ser Tyr Leu Lys Glu Glu Asp Glu Gly Lys Ile Asn
 690 695 700
 Tyr Val Trp Met Tyr Met Ile Glu Asn Asn Gln Ala Ser Ile Asp Asp
 705 710 715 720
 Ser Val Arg His Val Gln Thr Ile Ile Asn Val Lys Lys Gln Glu Phe
 725 730 735
 Ile Gln Arg Val Leu Ser Asp Gln His Cys Asn Leu Pro Lys Ser Phe
 740 745 750
 Lys Gln Leu His Phe Ser Cys Leu Lys Val Phe Asn Met Phe Phe Asn
 755 760 765
 Ser Ser Asn Ile Phe Asp Thr Asp Thr Asp Leu Leu Leu Asp Ile His
 770 775 780
 Glu Ala Phe Val Ser Pro Pro Gln Val Pro Lys Phe Lys Pro His Ile
 785 790 795 800
 Lys Pro Pro His Gln Leu Pro Ala Thr Leu Gln Pro Pro His Gln Pro
 805 810 815
 Gln Gln Ile Met Val Asn Lys Lys Lys Val Glu Met Val Tyr Lys Ser
 820 825 830
 Tyr His His Pro Phe Lys Val Phe Thr Leu Gln Lys Lys Gln Ser Ser
 835 840 845
 Gly His Gly Thr Met Asn Pro Arg Ala Ser Ile Leu Ala Gly Pro Asn
 850 855 860
 Ile Lys Leu Cys Phe Ser
 865 870



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/82, 9/88, 9/10, 9/00, 15/11, A01H 5/00	A3	(11) International Publication Number: WO 00/22150 (43) International Publication Date: 20 April 2000 (20.04.00)
(21) International Application Number: PCT/US99/23180 (22) International Filing Date: 5 October 1999 (05.10.99) (30) Priority Data: 09/172,339 14 October 1998 (14.10.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/172,339 (CON) Filed on 14 October 1998 (14.10.98) (71) Applicant (for all designated States except US): PIONEER HI-BRED INTERNATIONAL, INC. [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50319 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): MEYER, Terry, EuClaire [US/US]; 4338 101st Street, Urbandale, IA 50322 (US). YALPANI, Nasser [CA/US]; 6041 North Winwood Drive, Johnston, IA 50131 (US). (74) Agents: SPRUILL, W., Murray et al.; Alston & Bird LLP, P.O. Drawer 34009, Charlotte, NC 28234-4009 (US).	(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. (88) Date of publication of the international search report: 5 October 2000 (05.10.00)	

(54) Title: LIMONENE AND OTHER METABOLITES OF GERANYL PYROPHOSPHATE FOR INSECT CONTROL



(57) Abstract

Methods for manipulating metabolic pathway in plants, particularly those pathways involved in the biosynthesis of monoterpenes are provided. Methods are directed at transforming plants with one or more nucleotide sequences encoding the enzyme GPP synthase, and the monoterpene synthases limonene-, carveol and S-linalool synthase. Methods for creating or enhancing resistance to insects in plants by transforming plants with GPP- and/or monoterpene synthases, to generate plants producing monoterpenes in amount effective for resistance to insects are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/23180

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/82 C12N9/88 C12N9/10 C12N9/00 C12N15/11
A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 37102 A (PIONEER HI BRED INT) 28 November 1996 (1996-11-28) the whole document ---	1-11, 13-16, 20-30
X	WO 94 22304 A (PIONEER HI BRED INT) 13 October 1994 (1994-10-13) the whole document ---	1-7, 20-30
X	WO 95 11913 A (UNIV WASHINGTON) 4 May 1995 (1995-05-04) the whole document ---	1-7, 20-30
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

5 April 2000

Date of making of the international search report

25. 07. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Oderwald, H

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/23180

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>COLBY S M ET AL: "4S-LIMONENE SYNTHASE FROM THE OIL GLANDS OF SPEARMINT (MENTHA SPICATA)" JOURNAL OF BIOLOGICAL CHEMISTRY,US,AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, vol. 268, no. 31, 5 November 1993 (1993-11-05), pages 23016-23024, XP002914702 ISSN: 0021-9258 the whole document -----</p>	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 99/23180

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

SEE ADDITIONAL SHEET

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-11, 13-18, 20-30 - partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: {1-11, 13-18, 20-30 partially}

A method for manipulating a metabolic pathway in a plant cell comprising transforming a plant cell with a limonene synthase-encoding nucleotide sequence set forth in SEQ ID NO: 1 and coding for a protein set forth in SEQ ID NO: 2. A method for creating or enhancing resistance to insects in a plant, a method for producing limonene synthase in a plant cell, a transformed plant cell and plant having a manipulated metabolic pathway comprising said limonene synthase.

2. Claims: {1-30 partially}

same as invention 1 but comprising a GPP synthase as set forth in SEQ ID NO: 3 and 4.

3. Claims: {1-9, 11, 13-15, 17, 18, 20-30 partially}

same as in invention but comprising a carveol synthase as set forth in SEQ ID NO: 5 and 6.

4. Claims: {1-9, 12-15, 19-30 partially}

same as invention 1 but comprising an S-linalool synthase as set forth in SEQ ID NO: 7 and 8.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/23180

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9637102 A	28-11-1996	AT 185043 T	15-10-1999
		AU 5879496 A	11-12-1996
		BR 9609204 A	11-05-1999
		CA 2222023 A	28-11-1996
		DE 69604494 D	04-11-1999
		DE 69604494 T	05-01-2000
		EP 0836381 A	22-04-1998
		ES 2139361 T	01-02-2000

WO 9422304 A	13-10-1994	AT 185042 T	15-10-1999
		AU 6449794 A	24-10-1994
		CA 2159735 A	13-10-1994
		DE 69420957 D	04-11-1999
		DE 69420957 T	13-01-2000
		EP 0691812 A	17-01-1996

WO 9511913 A	04-05-1995	AU 8127994 A	22-05-1995
		CA 2118071 A	29-04-1995
		EP 0730603 A	11-09-1996
		US 5871988 A	16-02-1999

